

## 2. Correlations between Stellar Properties

### 2.1 Mass-luminosity relationship (ZG: 12.2; CO: 7.3)

- Most stars obey

$$L_s = \text{constant} \times M_s^\nu \quad 3 < \nu < 5$$



*Exercise 2.1:* Assuming a Salpeter IMF, show that most of the mass in stars in a galaxy is found in low-mass stars, while most of the stellar light in a galaxy comes from massive stars.

2.2 *Hertzsprung–Russell diagram* (ZG: 13-3; CO: 8.2) (plot of  $L_s$  vs.  $T_{\text{eff}}$ ): and *Colour–Magnitude Diagram* (e.g. plot of V vs. B-V) From diagrams for nearby stars of known distance we deduce:

1. About 90% of stars lie on the main sequence (broad band passing diagonally across the diagram)
2. Two groups are very much more luminous than MS stars (giants and supergiants)
3. One group is very much less luminous; these are the white dwarfs with  $R_s \ll R_\odot$  but  $M_s \sim M_\odot$ .

$\log g - \log T_{\text{eff}}$  diagram, determined from atmosphere models (does not require distance)

### Hertzsprung-Russell (Colour-Magnitude) Diagram



Hipparcos (1989 - 1993)




Fig. 2. Mass-luminosity relationship for the luminosity V stars from this paper (plus signs), from Popper (1980) (points) and from Heintze (1973) (broken line).




Fig. 3. The diagram  $\log g, \log T_{\text{eff}}$  for stars of different luminosities plotted from the data of Ta $\ddot{a}$

## 2.3 Cluster H-R Diagrams (ZG:13-3, 14-2; OG: 13.4)

- *Galactic or open clusters* – 10 to 1000 stars, not concentrated towards centre of cluster – found only in *disc of Galaxy*
- *Globular clusters* – massive spherical associations containing  $10^5$  or more stars, *spherically distributed* about centre of Galaxy, many at great distances from plane.
- All stars within a given cluster are effectively *equidistant* from us; we are probably seeing *homogeneous, coeval* groups of stars, and with the *same chemical composition*. We can construct *H-R diagrams* of apparent brightness against temperature.

### Main features of H-R diagrams:

#### 1. Globular clusters

- All have *main-sequence turn-offs* in similar positions and giant branches joining the main sequence at that point.
- All have *horizontal branches* running from near the top of the giant branch to the main sequence above the turn-off point.
- In many clusters *RR Lyrae stars* (of variable luminosity) occupy a region of the horizontal branch.

#### 2. Galactic clusters


- Considerable variation in the MS turn-off point; lowest in about the same position as that of globular clusters.
- Gap between MS and giant branch (*Hertzsprung gap*) in clusters with high turn-off point.

# STAR CLUSTERS




Open Cluster (Pleiades)


Globular Cluster (47 Tuc)



HST




47 Tucanae




Chandra (X-rays)


# Globular Cluster CM Diagrams



47 Tuc



M15



## 2.4 Chemical Composition of Stars (ZG: 13-3; CO: 9.4)

- We deduce the *photospheric composition* by studying *spectra*: information often incomplete and of doubtful precision.
- *Solar system* abundances: Reasonable agreement between analysis of solar spectrum and laboratory studies of meteorites (carbonaceous chondrites).
- *Normal stars* (vast majority): Similar composition to Sun and interstellar medium  
Typically: *Hydrogen 90% by number; Helium 10%; other elements (metals)  $\ll 1\%$*   
(by mass:  $X \simeq 0.70$ ,  $Y \simeq 0.28$ ,  $Z \simeq 0.02$ )
- *Globular cluster stars*: *Metal deficient* compared to Sun by factors of 10–1000,  
Hydrogen and helium normal

Assuming uniform initial composition for the Galaxy, we conclude that about 99% of metals must have been synthesized within stars.

**THIS IS THE PRIMARY EVIDENCE FOR  
NUCLEOSYNTHESIS DURING STELLAR EVOLUTION.**

## 2.5 STELLAR POPULATIONS (ZG: 14-3; CO: 13.4)

Population I: metallicity:  $Z \sim 0.02$  (i.e. solar), old and young stars, mainly in the Galactic disc, open clusters

Population II: metallicity:  $Z \sim 0.1 - 0.001 Z_{\odot}$ , old, high-velocity stars in the Galactic halo, globular clusters

Population III: hypothetical population of zero-metallicity stars (first generation of stars?), possibly with very different properties (massive, leading to relatively massive black holes?), may not exist as a major separate population (HE0107-5240, a low-mass star with  $Z \sim 10^{-7}$ : the first pop III star discovered?)

### *Stars with peculiar surface composition*

- Most stars seem to retain their initial surface composition as the centre evolves. A small number show anomalies, which can occur through:
  - 1) *mixing* of central material to the surface
  - 2) large scale *mass loss* of outer layers exposing interior (e.g. helium stars)
  - 3) *mass transfer* in a binary (e.g. barium stars)
  - 4) pollution with *supernova* material from a binary companion (e.g. Nova Sco)

### *Sub-stellar objects*

- *Brown Dwarfs*: star-like bodies with *masses too low* to create the central temperature required to *ignite fusion reactions* (i.e.  $M \lesssim 0.08 M_{\odot}$  from theory).
- *Planets*: *self-gravitating* objects formed in *disks* around stars (rocky planets [e.g. Earth], giant gas planets [e.g. Jupiter])

## *Summary II*

### Concepts:

- How does one determine *mass-luminosity relations*?
- The importance of the *Hertzsprung-Russell* and *Colour-Magnitude diagram*
- Basic properties of *open and globular clusters*
- The *chemical composition* of stars (metallicity)
- The different *stellar populations*
- Difference between *stars, brown dwarfs and planets*