

Oxford Physics
M. Phys. Major Option C1
Astrophysics



Cosmology & Galaxies Lecture Notes (2019-2020)

Julien Devriendt

DENYS WILKINSON BUILDING,
KEBLE ROAD, OXFORD OX1 3RH

`julien.devriendt@physics.ox.ac.uk`

JD—MAY 30, 2020

1 Disclaimer

Special thanks to Pedro Ferreira who kindly provided his Cosmology lectures notes and Adrienne Slyz who graciously shared the material of a chapter she wrote on the relation between fine-tuning and large scale structure (OUP, J. Silk et al editors). The first part of my lectures notes heavily draws from this source, and in particular covers the same material and adopts a very similar structure. I have also endeavoured to use the same notation as that of Steve Balbus' lectures on GR & Cosmology in the 3rd year course. Thus, although I present a brief overview of the homogeneous Universe model at the beginning of these lecture notes to make them more self-contained and refresh the memory of the reader concerning this notation, I assume that the material related to this topic is known.

These notes are, by and large, a direct transcription of what I write on the blackboard (with the exception of GR perturbation theory where I provide more intermediate steps in this manuscript). Thus they should be considered as an introduction to the subject rather than a definitive treatment. I perused many text books to write these notes, and here are my main sources of inspiration:

[L]: A. Liddle, *An Introduction to Modern Cosmology*, Wiley

[P]: J. Peacock, *Cosmological Physics*, CUP

[TP] T. Padmanabhan, *Large Scale Structure*, CUP

[D]: S. Dodelson, *Modern Cosmology*, Academic Press

[KT]: E. Kolb & M. Turner, *The Early Universe*, Addison Wesley.

[SG]: L.S. Sparke & J.S. Gallagher, *Galaxies in the Universe*, CUP

[BT]: J. J. Binney & S. Tremaine, *Galactic Dynamics*, PUP

[BM]: J. J. Binney & M. Merrifield, *Galactic Astronomy*, PUP

[W]: S. Weinberg, *Gravitation and Cosmology*, Wiley

In each section of these notes, I have suggested which of these books I believe is best suited to study the topic at hand in more depth.

Finally, I apologise in advance for any typo/error in these notes: please do not hesitate to contact me when you find one (including a departure of notation from the 3rd year course) so I can get the chance to fix them quickly. This will be much appreciated.

2 The Homogeneous Universe

As was established in the 3rd year GR & Cosmology course, the most general spacetime metric in an homogeneous and isotropic Universe, where one can define a uniform time is given by (see e.g. [W]),

chapter 13 for proof):

$$\begin{aligned}
 ds^2 &= g_{\mu\nu} dx^\mu dx^\nu \\
 &= (cdt \ dr \ d\theta \ d\phi) \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & a^2[t]/(1 - Kr^2) & 0 & 0 \\ 0 & 0 & a^2[t]r^2 & 0 \\ 0 & 0 & 0 & a^2[t]r^2 \sin^2 \theta \end{pmatrix} \begin{pmatrix} cdt \\ dr \\ d\theta \\ d\phi \end{pmatrix} \\
 &= -c^2 dt^2 + a^2[t] \left(\frac{dr^2}{1 - Kr^2} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \right) \tag{1}
 \end{aligned}$$

where $g_{\mu\nu}$ is the (diagonal) *covariant* metric tensor, c is the speed of light in vacuum, K is a constant indicative of the spatial curvature of the Universe and $a[t]$ is the time dependent expansion factor*. r, θ, ϕ define a *comoving* (spherical) coordinate system — i.e. a system in which an observer at rest moves along with the expansion of the Universe. This metric is called the Friedmann–Lemaître–Robertson–Walker (FLRW) metric, and we have adopted the space-like sign convention $(-, +, +, +)$ for the metric signature, as in the 3rd year course.

The same considerations about homogeneity and isotropy also imply that if we sit in the comoving frame in which the fluid[†] which fills the Universe is at rest[‡], its energy–momentum tensor can be written in a simple diagonal form as:

$$T_\mu^\nu = \begin{pmatrix} -\rho[t]c^2 & 0 & 0 & 0 \\ 0 & P[t] & 0 & 0 \\ 0 & 0 & P[t] & 0 \\ 0 & 0 & 0 & P[t] \end{pmatrix} \tag{2}$$

where $\rho[t]$ and $P[t]$ are the (time dependent) *average* density and pressure of the fluid. Indeed, starting from the general form of the energy-momentum tensor

$$T_{\mu\alpha} = (\rho[t] + P[t]/c^2)U_\mu U_\alpha + P[t]g_{\mu\alpha}$$

where $U_\mu = (-c, 0, 0, 0)$ is the *covariant* 4-velocity of the fluid at rest, we can multiply it by the *contravariant* (or inverse) metric tensor,

$$g^{\alpha\nu} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & (1 - Kr^2)/a^2[t] & 0 & 0 \\ 0 & 0 & 1/(a^2[t]r^2) & 0 \\ 0 & 0 & 0 & 1/(a^2[t]r^2 \sin^2 \theta) \end{pmatrix}$$

* $K = 0$ for a flat (Euclidean) universe, $K > 0$ for a closed (spherical) universe and $K < 0$ for an open (hyperbolic) one. In what follows, contrary to what was done in the 3rd year course, we will choose that $K^{-1/2}$ (= a in 3rd year course parlance), like r , has the dimensions of a length, so as to be able to normalise the *dimensionless* expansion factor (called $a[t]$ here instead of $R[t]$ in the 3rd year course) to unity at present, i.e. set $a_0 \equiv a[t_0] = 1$. This arbitrary choice takes advantage of our freedom to simultaneously rescale a, K and r without changing the geometry of spacetime. Unless otherwise mentioned, in these notes the subscript '0' indicates time dependent quantities evaluated at the present time t_0 .

[†]We only consider *ideal* fluids in these lectures. For a discussion of cases when this approximation breaks down, see e.g. [D].

[‡]Note that this choice is even possible for a fluid of photons, in spite of these latter travelling at the speed of light in any reference frame, as the velocity of the co-moving frame is simply the bulk velocity of the fluid.

to obtain

$$T_{\mu}^{\nu} = (\rho[t] + P[t]/c^2)U_{\mu}U^{\nu} + P[t]\delta_{\mu}^{\nu}. \quad (3)$$

Using the normalisation $g^{\mu\nu}U_{\mu}U_{\nu} = U_{\mu}U^{\mu} = U_0U^0 = -c^2$ then yields equation (2). The nature of the fluid then needs to be specified by an equation of state of the form $P = P[\rho]$.

Finally, Einstein's field equations derived in the 3rd year course — with the same convention (+,−,−) for the signs in front of the metric, Riemann tensor and energy-momentum tensor respectively — relate metric and energy momentum tensor, thus describing the interaction between space-time and matter:

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = -\frac{8\pi G}{c^4}T_{\mu\nu} + \Lambda g_{\mu\nu} \quad (4)$$

where $R_{\mu\nu}$ and R , the Ricci tensor and scalar, are functions of $g_{\mu\nu}$ and its first and second derivatives with respect to coordinates, G is the universal gravitational constant and Λ is the so-called cosmological constant. As you have seen in the 3rd year course, using the explicit forms of the FLRW metric and ideal fluid energy-momentum tensor given by equations (1) and (2), these field equations (4) yield the following pair of simple independent equations (R_{00} and $R_{ij}\delta_{ij}$ ($= R_{rr}$) terms respectively):

$$\left(\frac{da}{dt}\right)^2 = \frac{8\pi G a^2[t]}{3}\rho[t] + \frac{\Lambda a^2[t]c^2}{3} - Kc^2 \quad (5)$$

$$\frac{d^2a}{dt^2} = -\frac{4\pi G a[t]}{3}\left(\rho[t] + 3\frac{P[t]}{c^2}\right) + \frac{\Lambda a[t]c^2}{3} \quad (6)$$

called the Friedmann–Lemaître (FL) equations. Together with the equation of state of the fluid, these two equations constitute a closed system, i.e. they completely describe the dynamical evolution of an homogeneous and isotropic Universe.

It is convenient to define the *Hubble parameter* (a.k.a. the expansion rate of the Universe), $H[t] = \dot{a}[t]/a[t]$, where the dot over $a[t]$ stands for the derivative w.r.t. time, and a *critical density*

$$\rho_c[t] = \frac{3H^2[t]}{8\pi G}$$

obtained by setting $\Lambda = 0$ and $K = 0$ in the first FL equation (i.e. equation (5)). Finally, rather than the expansion factor, observers will talk about the cosmological *redshift*, $z \equiv a_0/a[t] - 1$, as it is the quantity which is directly observable. Given these definitions, one can rewrite equation (5) as:

$$\rho_c[z] = \rho[z] + \frac{\Lambda c^2}{8\pi G} - \frac{3Kc^2(1+z)^2}{8\pi G}$$

We can also compare all energy densities of the form $\rho_X[z]c^2$ — where the subscript $X = \{\gamma, M, V, K\}$ indicates contributions from relativistic matter, non-relativistic matter, vacuum energy and curvature respectively — to the critical energy density value $\rho_c[z]c^2$, and write the ratios as dimensionless

functions of redshift, $\Omega_X[z] = \rho_X[z]/\rho_c[z]$. We already have $\rho_V[z] = \Lambda c^2/(8\pi G)$ and $\rho_K[z] = -3Kc^2(1+z)^2/(8\pi G)$ in the previous equation, so we just need to separate $\rho[z]$ into a non-relativistic contribution $\rho_M[z]$ and a relativistic one $\rho_\gamma[z]$ to obtain:

$$\Omega_M[z] + \Omega_\gamma[z] + \Omega_V[z] + \Omega_K[z] = 1 \quad (7)$$

Note that only $\Omega_K[z]$ can be negative in this equation. Making the scalings with redshift explicit, i.e. $\rho_M[z] \propto (1+z)^3$, $\rho_\gamma[z] \propto (1+z)^4$, $\rho_V[z] = \text{constant}$, $\rho_K[z] \propto (1+z)^2$ and choosing as a reference the present day values, we can rewrite the equation governing the expansion rate of the Universe in the following convenient form:

$$H^2[z] = H_0^2 \left(\Omega_{M,0}(1+z)^3 + \Omega_{\gamma,0}(1+z)^4 + \Omega_{V,0} + \Omega_{K,0}(1+z)^2 \right) \quad (8)$$

Current best estimates for these are $\Omega_{M,0} = 0.317$, $\Omega_{V,0} = 0.683$, $\Omega_{K,0} = 0.0$, $\Omega_{\gamma,0} = 9 \times 10^{-5}$, and $H_0 = 67.2 \text{ km/s/Mpc}$ (c.f. Planck collaboration results 2018)[§].

Going back to equation (1), one notices that by defining a new time variable η , called the *conformal* time, as $d\eta \equiv dt/a[t]$, it is possible to factor the expansion of the Universe out of the spacetime interval ds^2 , which then reads:

$$ds^2 = a^2[\eta] \left(-c^2 d\eta^2 + \frac{dr^2}{1 - Kr^2} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \right). \quad (9)$$

η is the amount of time it would take a photon to travel from where an observer is located to the largest distance it can observe, provided the Universe stopped expanding. Which leads us to define the important notion of *comoving*, or *particle horizon*, $R_h[t]$, as the maximum distance a massless particle can have travelled since the Big Bang:

$$R_h[t] = c\eta = c \int_0^t \frac{dt'}{a[t']} = R_h[z] = c \int_z^\infty \frac{dz'}{H[z']} \quad (10)$$

Note that since the *proper* distance is equal to the comoving distance times the expansion factor (i.e. these two distances are equal today, at $z = 0$, because by convention $a_0 = 1$), the proper distance to the particle horizon at any redshift is simply $R_h[z]/(1+z)$.

3 Newtonian Evolution of Cosmological Perturbations

So far, we have built our understanding of the expanding Universe on an extraordinary simplifying assumption — that at any given time, it looks the same everywhere and any direction. This assumption is underpinned by observations of the cosmic microwave background radiation which we find to be uniform to within one part in a hundred thousand. Yet we also know that the Universe cannot be

[§]In the remainder of these notes we will drop the '0' subscript and write $\Omega_i = \Omega_{i,0}$ i.e. assume these parameters have their present day values, unless the situation is ambiguous. Note that the Ω_V contribution in equation (8) can also be generalised to a more complex form of 'dark energy', with a different equation of state.

perfectly smooth. The latest observational surveys have identified large scale structures in abundance: galaxies grouped together in clusters, filaments and walls or separated by large empty voids that can span hundreds of millions of light years. Indeed, the mere existence of galaxies, stars and planets indicate that the Universe looks more and more anisotropic as scales become smaller and smaller.

Therefore, if we are to understand the formation of large scale structure, we must go one step beyond the homogeneous Universe and derive the differential equations which govern the growth of small inhomogeneities, i.e. allow for spatial variations in the evolution of energy density, pressure, and gravity. In this section, we will use Newtonian gravity when doing so. However, before we go any further, it is important to make the following disclaimer, to clarify the situation: *it is impossible to study cosmology using Newtonian gravity in a mathematically rigorous and self-consistent way, for all type of fluids and at all perturbation wavelengths.* Having said that, as we will see in the following section, Newtonian gravity does provide us with the qualitative and quantitative behaviour of perturbations that we would find in a proper, general relativistic treatment, at least in the case of massive, non-relativistic particles.

More specifically, let us consider the case where matter is an ideal, non-relativistic self-gravitating fluid. At the epoch where it is the dominant component* the equations governing its dynamics are:

$$\frac{\partial \rho}{\partial t}[\vec{r}, t] + \frac{\partial}{\partial \vec{r}}(\rho \vec{u})[\vec{r}, t] = 0, \quad (11)$$

where \vec{r} is the position of a fluid element in physical (proper) coordinates and \vec{u} its corresponding velocity. Equation (11) represents mass (energy) conservation. We also have conservation of momentum (Euler's equations)

$$\frac{\partial \vec{u}}{\partial t}[\vec{r}, t] + \left(\vec{u} \frac{\partial}{\partial \vec{r}} \right) \vec{u}[\vec{r}, t] = -\frac{\partial}{\partial \vec{r}} \Psi[\vec{r}, t] - \frac{1}{\rho} \frac{\partial}{\partial \vec{r}} P[\vec{r}, t], \quad (12)$$

and finally, Poisson's equation for the gravitational potential Ψ which writes[†]:

$$\left(\frac{\partial}{\partial \vec{r}} \right)^2 \Psi[\vec{r}, t] = 4\pi G \rho[\vec{r}, t]. \quad (13)$$

Rigorously speaking, these equations are only valid on scales which are small compared to the particle horizon previously defined, and when the fluid is far away enough from a black hole.

One can more conveniently re-write these equations in a comoving coordinate system defined by $\vec{x} = \vec{r}/a[t]$. In such a system, we have $\vec{u} = \dot{a}\vec{x} + \vec{v}[\vec{x}, t]$ where $\dot{a}\vec{x}$ is called the *Hubble flow* as it is the

*This is less restrictive than it may seem, given the values of the cosmological parameters quoted in the previous section. According to these our Universe becomes matter dominated when $\Omega_M[z] = \Omega_\gamma[z]$, i.e at $z \sim 3400$ and stops being so when $\Omega_V[z] = \Omega_M[z]$, i.e. at $z \sim 0.3$. This spans a time window of ~ 10.2 Gyr, to be compared to the age of the Universe, ~ 13.7 Gyr.

†If one wanted to include the cosmological constant Λ it would appear on the RHS of this equation with a negative sign.

velocity contributed by the expansion of the Universe and $\vec{v}[\vec{x}, t]$ is the *peculiar* velocity of the fluid element. This leads to[‡]:

$$\frac{\partial \rho}{\partial t}[\vec{x}, t] + \frac{3\dot{a}}{a}\rho[\vec{x}, t] + \frac{1}{a}\frac{\partial}{\partial \vec{x}}(\rho\vec{v})[\vec{x}, t] = 0 \quad (14)$$

$$\frac{\partial \vec{v}}{\partial t}[\vec{x}, t] + \frac{1}{a}\left(\vec{v}\frac{\partial}{\partial \vec{x}}\right)\vec{v}[\vec{x}, t] + \frac{\dot{a}}{a}\vec{v}[\vec{x}, t] = -\frac{1}{a}\frac{\partial}{\partial \vec{x}}\Phi[\vec{x}, t] - \frac{1}{a\rho}\frac{\partial}{\partial \vec{x}}P[\vec{x}, t] \quad (15)$$

$$\left(\frac{1}{a}\frac{\partial}{\partial \vec{x}}\right)^2\Phi[\vec{x}, t] = 4\pi G\rho[\vec{x}, t] + 3\frac{\ddot{a}}{a} \quad (16)$$

where we have defined a new gravitational potential $\Phi = \Psi + \frac{1}{2}a\ddot{a}\vec{x}^2$ to cast equation (15) in a simpler form.

In the limit where the fluid is only slightly perturbed compared to the homogeneous and isotropic background (which we know is a valid assumption at the time when the cosmic microwave background is observed, around $z \sim 1100$) we can write $\rho[\vec{x}, t] = \bar{\rho}[t](1 + \delta[\vec{x}, t])$ with the density contrast $\delta[\vec{x}, t] \equiv (\rho[\vec{x}, t] - \bar{\rho}[t])/\bar{\rho}[t] \ll 1$, $\vec{v}[\vec{x}, t] \ll \dot{a}\vec{x}$, $P[\vec{x}, t] = \bar{P}[t] + \delta P[\vec{x}, t] = \bar{P}[t] + (\partial P/\partial \rho)_S \delta\rho[\vec{x}, t] = \bar{P}[t] + c_s^2\bar{\rho}\delta[\vec{x}, t]$ for adiabatic perturbations with associated sound speed c_s , $\Phi[\vec{x}, t] \ll \frac{1}{2}a\ddot{a}\vec{x}^2$ and the bar above a variable indicates the homogeneous background average.

Linearising equations (14) (15) and (16), i.e. using the FL equations to cancel the 0th order terms and keeping only the first order terms in all the perturbed variables (which we leave as an exercise), one obtains:

$$\frac{\partial \delta}{\partial t}[\vec{x}, t] + \frac{1}{a}\frac{\partial \vec{v}}{\partial \vec{x}}[\vec{x}, t] = 0 \quad (17)$$

$$\frac{\partial \vec{v}}{\partial t}[\vec{x}, t] + \frac{\dot{a}}{a}\vec{v}[\vec{x}, t] = -\frac{1}{a}\frac{\partial}{\partial \vec{x}}\Phi[\vec{x}, t] - \frac{c_s^2}{a}\frac{\partial}{\partial \vec{x}}\delta[\vec{x}, t] \quad (18)$$

$$\nabla_{\vec{x}}^2\Phi[\vec{x}, t] = 4\pi G\bar{\rho}a^2\delta[\vec{x}, t] \quad (19)$$

Finally, taking the gradient of the perturbed Euler equation (18) and using equations (17) and (19) to get rid of $\nabla_{\vec{x}}\vec{v}$ and $\nabla_{\vec{x}}^2\Phi$ respectively, yields the following 2nd order linear partial differential equation for δ :

$$\ddot{\delta}[\vec{x}, t] + 2H\dot{\delta}[\vec{x}, t] - \frac{c_s^2}{a^2}\nabla_{\vec{x}}^2\delta[\vec{x}, t] = 4\pi G\bar{\rho}\delta[\vec{x}, t]. \quad (20)$$

This equation is more easily studied and solved in Fourier space. Taking its Fourier transform[§],

[‡]Being careful about the variable changes:

$$\left(\frac{\partial}{\partial t}\right)_{\vec{r}} = \left(\frac{\partial}{\partial t}\right)_{\vec{x}} - \frac{\dot{a}}{a}\left(\vec{x}\cdot\frac{\partial}{\partial \vec{x}}\right)_t \quad \& \quad \left(\frac{\partial}{\partial \vec{r}}\right)_t = \frac{1}{a}\left(\frac{\partial}{\partial \vec{x}}\right)_t$$

[§]The Fourier transform of δ is defined as

$$\delta[\vec{x}, t] = \frac{1}{(2\pi)^3} \int d^3k \delta_{\mathbf{k}} \exp(-i\vec{k}\cdot\vec{x})$$

equation (20) becomes

$$\ddot{\delta}_{\mathbf{k}} + 2H\dot{\delta}_{\mathbf{k}} + \left(\frac{c_s^2}{a^2} k^2 - 4\pi G\bar{\rho} \right) \delta_{\mathbf{k}} = 0, \quad (21)$$

or

$$\delta_{\mathbf{k}}'' + \mathcal{H}\delta_{\mathbf{k}}' + (c_s^2 k^2 - 4\pi G a^2 \bar{\rho}) \delta_{\mathbf{k}} = 0 \quad (22)$$

using the conformal variables defined in equation (9), and replacing the partial derivative w.r.t. conformal time, $\partial/\partial\eta \rightarrow '.$ The *conformal* Hubble parameter is thus $\mathcal{H} \equiv a'/a = aH.$ This is arguably the most important equation of these Cosmology lectures, as we will come back to it time and again to understand the growth of structures in the Universe, even in the non-linear regime relevant to galaxy formation!

4 Relativistic Cosmological Perturbation Theory

The Newtonian analysis developed in the previous section clearly has limitations. In particular, it fails for perturbations with sizes comparable or larger than the particle horizon, and for fully relativistic fluids at both large and short wavelengths, as the fluid pressure also contributes significantly to the active gravitational mass in this case, unlike in the Newtonian case where only the inertial mass matters.

Unfortunately, the physical interpretation of the results obtained is less transparent in General Relativity (GR) than in the Newtonian theory of gravitation. The main problem arises from the freedom in the choice of coordinates that GR allows to describe perturbations. Indeed, in contrast to the homogeneous and isotropic Universe where the preferable coordinate system is fixed by the symmetry properties there is no obvious choice as soon as we introduce perturbations. Freedom in the coordinate choice, called *gauge freedom*, generally leads to the appearance of fictitious perturbation modes which only reflect the properties of the coordinate system used.

To illustrate this point, let us consider the unperturbed homogeneous and isotropic Universe, where $\rho[\vec{x}, t] = \bar{\rho}[t],$ and define a new time coordinate, t' related to t via $t' = t + \delta t[\vec{x}, t].$ The mass (energy) density in this new coordinate system, $\rho'[\vec{x}, t'] \equiv \bar{\rho}[t[\vec{x}, t']],$ when evaluated on the hypersurface $t' = C^{\text{st}},$ will in general depend on the spatial coordinate $\vec{x}.$ Assuming $\delta t \ll t$ we can then write $\bar{\rho}[t] = \bar{\rho}[t' - \delta t[\vec{x}, t]] \simeq \bar{\rho}[t'] - \dot{\bar{\rho}}\delta t[\vec{x}, t] \equiv \bar{\rho}[t'] + \delta\bar{\rho}[\vec{x}, t'],$ where the (fictitious) linear perturbation term $\delta\bar{\rho}[\vec{x}, t']$ is entirely due to our choice of a new "disturbed" time coordinate!

In the same way that fictitious terms can appear, real perturbation terms may disappear depending on our choice of coordinate system. There exist two (related) ways to overcome this problem. The first one is to calculate only combinations of metric perturbations which are gauge invariant, i.e. that are identical under any change of coordinate system. These combinations are called the *Bardeen*

variables. Alternatively, we can fix the gauge, and keep track of *all* perturbations, metric *and* matter. It is this approach that we will follow here, adopting the convenient Newtonian gauge, which is uniquely defined for perturbations that decay at spatial infinity* and commonly used to study the formation and evolution of large scale structure and cosmic microwave background anisotropies.

We will further restrict ourselves to *scalar* perturbations of the metric and associated energy density inhomogeneities, as these are the most relevant for structure formation, which is the main topic of these lectures†. Finally to avoid overly complicated calculations, but also because the measured values of the cosmological parameters given in the previous section indicate that our Universe satisfies this constraint to a high degree of accuracy, we will restrict ourselves to a flat FLRW spacetime.

More specifically, this means that our starting point will be the conformal metric of equation (9), with $K = 0$, cartesian spatial coordinates instead of spherical coordinates as spherical symmetry is broken anyways when perturbations are present. We then perturb this metric using a single scalar component Φ , such that‡:

$$ds^2 = a^2[\eta] \left(- \left(1 + \frac{2\Phi}{c^2} \right) c^2 d\eta^2 + \left(1 - \frac{2\Phi}{c^2} \right) \delta_{ij} dx^i dx^j \right). \quad (23)$$

You should be familiar with the Newtonian limit where you set $a[\eta] = 1$ and Φ is the standard Newtonian gravitational potential with the same definition than in the previous section (hence the name of (conformal) Newtonian gauge). As is customary, wherever we use latin indices, they run over the spatial part only, whereas greek indices span the whole of spacetime. We can view this metric as a 0th order part (which is just the usual FLRW metric) and a linearly perturbed part, i.e. as $g_{\mu\nu} = \bar{g}_{\mu\nu} + \delta g_{\mu\nu}$ with $\bar{g}_{\mu\nu}$ the usual metric of the homogeneous background Universe.

From there, we need to work out the linearly perturbed Einstein field equations which, if we ignore the cosmological constant term on the RHS§, will formally look like:

$$\delta G_{\mu\nu} = -\frac{8\pi G}{c^4} \delta T_{\mu\nu}, \quad (24)$$

*see [D] for a discussion of what specific gauge choices entail.

†Once again we refer to [D] for a discussion of the scalar-vector-tensor decomposition of metric perturbations, but note that, in any case, in the linear regime which we will explore in these lectures, these different perturbation modes evolve independently of one another, so that our results for scalar perturbations will hold regardless of whether vector or tensor modes are present.

‡A couple of remarks. Firstly, we use this form because it makes direct contact with the gauge invariant Bardeen variables previously discussed. Second, the scalar perturbations of the time part and spatial part of the metric have a priori no reason to be the same, which is why if you look at Cosmology text books, you will see two different variables Ψ and Φ used. In writing the metric as we do here, we are ignoring any source of anisotropic stress, in which case $\Psi = \Phi$. Since anisotropic stress generally is a second order effect, ignoring it is a reasonable approximation in linear perturbation theory.

§It is easy to add it back in at the very end, so we leave it as an exercise, especially since we ignored its presence in the Newtonian theory as well in the previous section and we wish to make the results of the two sections as easily comparable as possible.

where we have written the Einstein tensor $G_{\mu\nu}$ as the usual linear combination of the Ricci tensor $R_{\mu\nu}$ and the metric tensor times the Ricci scalar R :

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R.$$

Let us start with evaluating the LHS of equation (24). To do so, the first step is to calculate the connection coefficients:

$$\Gamma_{\nu\rho}^{\mu} = \frac{1}{2}g^{\mu\alpha}(\partial_{\nu}g_{\alpha\rho} + \partial_{\rho}g_{\alpha\nu} - \partial_{\alpha}g_{\nu\rho}). \quad (25)$$

We thus write the co- and contra-variant conformal metric of equation (23):

$$g_{\alpha\beta} = a^2[\eta] \begin{pmatrix} -(1 + 2\Phi/c^2) & 0 \\ 0 & (1 - 2\Phi/c^2)\delta_{ij} \end{pmatrix}$$

and

$$g^{\alpha\beta} = \frac{1}{a^2[\eta]} \begin{pmatrix} -(1 - 2\Phi/c^2) & 0 \\ 0 & (1 + 2\Phi/c^2)\delta^{ij} \end{pmatrix}$$

respectively, where we used the fact that for $\epsilon \ll 1$, $(1 + \epsilon)^{-1} \simeq 1 - \epsilon$. We then substitute these in the definition of the connection coefficients of equation (25) to obtain[¶]:

$$\begin{aligned} \Gamma_{00}^0 &= \frac{1}{2}g^{00}\partial_0g_{00} = \frac{-1 + 2\Phi/c^2}{2a^2}\partial_0(-a^2(1 + 2\Phi/c^2)) &= \frac{\mathcal{H}}{c} + \frac{\Phi'}{c^3} + \mathcal{O}\left(\frac{\Phi^2}{c^4}\right) \\ \Gamma_{0i}^0 &= \frac{1}{2}g^{00}\partial_i g_{00} = \frac{-1 + 2\Phi/c^2}{2a^2}\partial_i(-a^2(1 + 2\Phi/c^2)) &= \frac{\partial_i\Phi}{c^2} \\ \Gamma_{00}^i &= \frac{1}{2}g^{ij}(-\partial_j g_{00}) = \frac{(1 + 2\Phi/c^2)\delta^{ij}}{2a^2}\partial_j(-a^2(1 + 2\Phi/c^2)) &= \frac{\partial^i\Phi}{c^2} \\ \Gamma_{ij}^0 &= \frac{1}{2}g^{00}(-\partial_0 g_{ij}) = \frac{1 - 2\Phi/c^2}{2a^2}\partial_0(a^2(1 - 2\Phi/c^2)\delta_{ij}) &= \left(\frac{\mathcal{H}}{c} - 4\mathcal{H}\frac{\Phi}{c^3} - \frac{\Phi'}{c^3}\right)\delta_{ij} \\ \Gamma_{j0}^i &= \frac{1}{2}g^{ik}(\partial_0 g_{kj}) = \frac{(1 + 2\Phi/c^2)\delta^{ik}}{2a^2}\partial_0(a^2(1 - 2\Phi/c^2)\delta_{kj}) &= \left(\frac{\mathcal{H}}{c} - \frac{\Phi'}{c^3}\right)\delta_j^i \\ \Gamma_{jk}^i &= \frac{1}{2}g^{il}(\partial_j g_{lk} + \partial_k g_{lj} - \partial_l g_{jk}) &= -\frac{\partial_j\Phi}{c^2}\delta_k^i - \frac{\partial_k\Phi}{c^2}\delta_j^i + \frac{\partial_l\Phi}{c^2}\delta_{jk}\delta^{il}. \end{aligned}$$

where we have, by analogy with equation (22), used the following notation for the partial derivative w.r.t. conformal time: $c\partial_0 \rightarrow \partial/\partial\eta \rightarrow '$. Notice that the expressions for Γ_{00}^0 , Γ_{ij}^0 and Γ_{j0}^i have a 0th order bit (the \mathcal{H} terms) and a first order bit – the 0th order bit you have already derived when working out the FL equations (5) from scratch in the 3rd year GR & Cosmology course.

[¶]Remember we neglect all quadratic or higher order terms in small quantities – which we denote generically by $\mathcal{O}(\Phi^2/c^4)$ in the first calculation, but drop altogether in the remainder of these lecture notes.

Now we turn to combining the connection coefficients together to obtain the Ricci tensor:

$$R_{\mu\nu} = \partial_\nu \Gamma_{\mu\alpha}^\alpha - \partial_\alpha \Gamma_{\mu\nu}^\alpha + \Gamma_{\mu\alpha}^\beta \Gamma_{\nu\beta}^\alpha - \Gamma_{\mu\nu}^\beta \Gamma_{\alpha\beta}^\alpha.$$

Starting with the time component of the tensor

$$R_{00} = \partial_0 \Gamma_{0\alpha}^\alpha - \partial_\alpha \Gamma_{00}^\alpha + \Gamma_{0\alpha}^\beta \Gamma_{0\beta}^\alpha - \Gamma_{00}^\beta \Gamma_{\alpha\beta}^\alpha,$$

we notice that if $\alpha = 0$, each pair of terms cancels, so we only need to sum over $\alpha = i$. We are then left with four terms:

$$\begin{aligned} \partial_0 \Gamma_{0i}^i &= \partial_0 (\mathcal{H}/c - \Phi'/c^3) \delta_i^i &= 3 \frac{\mathcal{H}'}{c^2} - 3 \frac{\Phi''}{c^4} \\ \partial_i \Gamma_{00}^i &= \partial_i \partial^i \Phi / c^2 &= \frac{\nabla^2 \Phi}{c^2} \\ \Gamma_{0i}^\beta \Gamma_{0\beta}^i &= \cancel{\partial_i \Phi \partial^i \Phi / c^4} + (\mathcal{H}/c - \Phi'/c^3) \delta_i^j (\mathcal{H}/c - \Phi'/c^3) \delta_j^i &= 3 \frac{\mathcal{H}^2}{c^2} - 6\mathcal{H} \frac{\Phi'}{c^4} \\ \Gamma_{00}^\beta \Gamma_{i\beta}^i &= (\mathcal{H}/c + \Phi'/c^3) (\mathcal{H}/c - \Phi'/c^3) \delta_i^i &= 3 \frac{\mathcal{H}^2}{c^2}. \end{aligned}$$

So we find, keeping only up to first order terms in the perturbed potential, Φ :

$$R_{00} = 3 \frac{\mathcal{H}'}{c^2} - 3 \frac{\Phi''}{c^4} - \frac{\nabla^2 \Phi}{c^2} - 6\mathcal{H} \frac{\Phi'}{c^4}.$$

Note that, once again, it splits into a 0^{th} order bit (first term) and a linear bit (last 3 terms). Repeating this procedure for the mixed components of the Ricci tensor, one obtains:

$$R_{0i} = \partial_i \Gamma_{0\alpha}^\alpha - \partial_\alpha \Gamma_{0i}^\alpha + \Gamma_{0\alpha}^\beta \Gamma_{i\beta}^\alpha - \Gamma_{0i}^\beta \Gamma_{\alpha\beta}^\alpha,$$

and splitting the index α into time and spatial indices, we now have 8 terms to calculate:

$$\begin{aligned} \partial_i \Gamma_{00}^0 &= \partial_i (\mathcal{H}/c + \Phi'/c^3) &= \frac{\partial_i \Phi'}{c^3} \\ \partial_0 \Gamma_{0i}^0 &= \partial_0 \partial_i \Phi / c^2 &= \frac{\partial_i \Phi'}{c^3} \\ \Gamma_{00}^\beta \Gamma_{i\beta}^0 &= (\mathcal{H}/c + \Phi'/c^3) \partial_i \Phi / c^2 + \partial^j \Phi (\mathcal{H}/c - 4\mathcal{H}\Phi/c^3 - \Phi'/c^3) \delta_{ij} / c^2 &= 2\mathcal{H} \frac{\partial_i \Phi}{c^3} \\ \Gamma_{0i}^\beta \Gamma_{0\beta}^0 &= \partial_i \Phi (\mathcal{H}/c + \Phi'/c^3) / c^2 + (\mathcal{H}/c - \Phi'/c^3) \partial_i \Phi / c^2 &= 2\mathcal{H} \frac{\partial_i \Phi}{c^3} \\ \partial_i \Gamma_{0k}^k &= \partial_i (\mathcal{H}/c - \Phi'/c^3) \delta_k^k &= -3 \frac{\partial_i \Phi'}{c^3} \\ \partial_k \Gamma_{0i}^k &= \partial_k (\mathcal{H}/c - \Phi'/c^3) \delta_i^k &= -\frac{\partial_i \Phi'}{c^3} \\ \Gamma_{0k}^\beta \Gamma_{i\beta}^k &= \partial_i \Phi (\mathcal{H}/c - \Phi'/c^3) / c^2 + (\mathcal{H}/c - \Phi'/c^3) (-\delta_k^k \partial_i \Phi - \delta_i^j \partial_j \Phi + \delta_{ik} \delta^{kl} \partial_l \Phi) / c^2 &= -2\mathcal{H} \frac{\partial_i \Phi}{c^3} \\ \Gamma_{0i}^\beta \Gamma_{k\beta}^k &= \partial_i \Phi (\mathcal{H}/c - \Phi'/c^3) \delta_k^k / c^2 + (\mathcal{H}/c - \Phi'/c^3) \delta_i^j (-\delta_j^k \partial_k \Phi - \delta_k^k \partial_j \Phi + \delta_j^l \partial_l \Phi) / c^2 &= 0. \end{aligned}$$

So we simply have:

$$R_{0i} = -2\mathcal{H}\frac{\partial_i\Phi}{c^3} - 2\frac{\partial_i\Phi'}{c^3}.$$

Finally, for the spatial components of the Ricci tensor:

$$R_{ij} = \partial_j\Gamma_{i\alpha}^\alpha - \partial_\alpha\Gamma_{ij}^\alpha + \Gamma_{i\alpha}^\beta\Gamma_{j\beta}^\alpha - \Gamma_{ij}^\beta\Gamma_{\alpha\beta}^\alpha,$$

the same splitting of the index α yields the following 8 terms:

$$\begin{aligned} \partial_j\Gamma_{i0}^0 &= \partial_j\partial_i\Phi/c^2 &= \frac{\partial_i\partial_j\Phi}{c^2} \\ \partial_0\Gamma_{ij}^0 &= \partial_0(\mathcal{H}/c - 4\mathcal{H}\Phi/c^3 - \Phi'/c^3)\delta_{ij} &= \left(\frac{\mathcal{H}'}{c^2} - 4\mathcal{H}'\frac{\Phi}{c^4} - 4\mathcal{H}\frac{\Phi'}{c^4} - \frac{\Phi''}{c^4}\right)\delta_{ij} \\ \Gamma_{i0}^\beta\Gamma_{j\beta}^0 &= (\mathcal{H}/c - \Phi'/c^3)(\mathcal{H}/c - 4\mathcal{H}\Phi/c^3 - \Phi'/c^3)\delta_{ij} &= \left(\frac{\mathcal{H}^2}{c^2} - 4\mathcal{H}^2\frac{\Phi}{c^4} - 2\mathcal{H}\frac{\Phi'}{c^4}\right)\delta_{ij} \\ \Gamma_{ij}^\beta\Gamma_{0\beta}^0 &= (\mathcal{H}/c - 4\mathcal{H}\Phi/c^3 - \Phi'/c^3)(\mathcal{H}/c + \Phi'/c^3)\delta_{ij} &= \left(\frac{\mathcal{H}^2}{c^2} - 4\mathcal{H}^2\frac{\Phi}{c^4}\right)\delta_{ij} \\ \partial_j\Gamma_{ik}^k &= \partial_j(-\delta_k^k\partial_i\Phi - \delta_i^k\partial_k\Phi + \delta_{ik}\delta^{kl}\partial_l\Phi)/c^2 &= -3\frac{\partial_i\partial_j\Phi}{c^2} \\ \partial_k\Gamma_{ij}^k &= \partial_k(-\delta_i^k\partial_j\Phi - \delta_j^k\partial_i\Phi + \delta_{ji}\delta^{kl}\partial_l\Phi)/c^2 &= -2\frac{\partial_i\partial_j\Phi}{c^2} + \frac{\nabla^2\Phi}{c^2}\delta_{ij} \\ \Gamma_{ik}^\beta\Gamma_{j\beta}^k &= (\mathcal{H}/c - 4\mathcal{H}\Phi/c^3 - \Phi'/c^3)(\mathcal{H}/c - \Phi'/c^3)\delta_{ij} &= \left(\frac{\mathcal{H}^2}{c^2} - 4\mathcal{H}^2\frac{\Phi}{c^4} - 2\mathcal{H}\frac{\Phi'}{c^4}\right)\delta_{ij} \\ \Gamma_{ij}^\beta\Gamma_{k\beta}^k &= (\mathcal{H}/c - 4\mathcal{H}\Phi/c^3 - \Phi'/c^3)(\mathcal{H}/c - \Phi'/c^3)\delta_k^k\delta_{ij} = 3\left(\frac{\mathcal{H}^2}{c^2} - 4\mathcal{H}^2\frac{\Phi}{c^4} - 2\mathcal{H}\frac{\Phi'}{c^4}\right)\delta_{ij} \end{aligned}$$

which we collect to write:

$$R_{ij} = \left(-\frac{\mathcal{H}'}{c^2} - 2\frac{\mathcal{H}^2}{c^2} + 8\mathcal{H}^2\frac{\Phi}{c^4} + 4\mathcal{H}'\frac{\Phi}{c^4} + 6\mathcal{H}\frac{\Phi'}{c^4} + \frac{\Phi''}{c^4} - \frac{\nabla^2\Phi}{c^2}\right)\delta_{ij}.$$

The Ricci scalar is by definition:

$$R \equiv g^{\mu\nu}R_{\mu\nu} = g^{00}R_{00} + 2g^{0i}R_{0i} + g^{ij}R_{ij} = g^{00}R_{00} + g^{ij}R_{ij},$$

as $g^{0i} = 0$. Using the expressions previously derived for R_{00} and R_{ij} , we get:

$$\begin{aligned} Ra^2 &= -(1 - 2\Phi/c^2)(3\mathcal{H}'/c^2 - 3\Phi''/c^4 - \nabla^2\Phi/c^2 - 6\mathcal{H}\Phi'/c^4) + (1 + 2\Phi/c^2)\delta^{ij} \\ &\quad \times (-\mathcal{H}'/c^2 - 2\mathcal{H}^2/c^2 + 8\mathcal{H}^2\Phi/c^4 + 4\mathcal{H}'\Phi/c^4 + 6\mathcal{H}\Phi'/c^4 + \Phi''/c^4 - \nabla^2\Phi/c^2)\delta_{ij} \\ &= -6\frac{\mathcal{H}'}{c^2} - 6\frac{\mathcal{H}^2}{c^2} + 6\frac{\Phi''}{c^4} - 2\frac{\nabla^2\Phi}{c^2} + 24\mathcal{H}\frac{\Phi'}{c^4} + 12\mathcal{H}'\frac{\Phi}{c^4} + 12\mathcal{H}^2\frac{\Phi}{c^4}. \end{aligned}$$

We now have all the ingredients to work out the different components of the Einstein tensor. They are, starting with the time component:

$$\begin{aligned}
G_{00} &\equiv R_{00} - \frac{1}{2}g_{00}R \\
&= 3\mathcal{H}'/c^2 - 3\Phi''/c^4 - \nabla^2\Phi/c^2 - 6\mathcal{H}\Phi'/c^4 + \frac{1}{2}(1 + 2\Phi/c^2)(-6\mathcal{H}'/c^2 - 6\mathcal{H}^2/c^2 + 6\Phi''/c^4 \\
&\quad - 2\nabla^2\Phi/c^2 + 24\mathcal{H}\Phi'/c^4 + 12\mathcal{H}'\Phi/c^4 + 12\mathcal{H}^2\Phi/c^4) \\
&= -3\frac{\mathcal{H}^2}{c^2} - 2\frac{\nabla^2\Phi}{c^2} + 6\mathcal{H}\frac{\Phi'}{c^4}.
\end{aligned} \tag{26}$$

Once again, the first term on the RHS is the 0th order term you have derived in the 3rd year GR & Cosmology course, which yields (simply by dividing by -3 and remembering that we assumed a flat Universe, i.e. $K = 0$, without a cosmological constant, i.e. $\Lambda = 0$) the LHS term of the FL equation (5). The mixed space-time component is simply R_{0i} , since $g_{0i} = 0$, so:

$$G_{0i} = -2\mathcal{H}\frac{\partial_i\Phi}{c^3} - 2\frac{\partial_i\Phi'}{c^3}, \tag{27}$$

with no 0th order term, as expected from the FL equations for the homogeneous Universe! Finally, the spatial component reads:

$$\begin{aligned}
G_{ij} &\equiv R_{ij} - \frac{1}{2}g_{ij}R \\
&= (-\mathcal{H}'/c^2 + 4\mathcal{H}'\Phi/c^4 + 6\mathcal{H}\Phi'/c^4 + \Phi''/c^4 - \nabla^2\Phi/c^2 - 2\mathcal{H}^2/c^2 + 8\mathcal{H}^2\Phi/c^4) \delta_{ij} \\
&\quad - \frac{1}{2}(1 - 2\Phi/c^2)(-6\mathcal{H}'/c^2 - 6\mathcal{H}^2/c^2 + 6\Phi''/c^4 - 2\nabla^2\Phi/c^2 + 24\mathcal{H}\Phi'/c^4 + 12\mathcal{H}'\Phi/c^4 \\
&\quad + 12\mathcal{H}^2\Phi/c^4) \delta_{ij} \\
&= \left(2\frac{\mathcal{H}'}{c^2} + \frac{\mathcal{H}^2}{c^2} - 8\mathcal{H}'\frac{\Phi}{c^4} - 6\mathcal{H}\frac{\Phi'}{c^4} - 2\frac{\Phi''}{c^4} - 4\mathcal{H}^2\frac{\Phi}{c^4} \right) \delta_{ij}.
\end{aligned} \tag{28}$$

with the 0th order term providing two of the terms in the FL equation (6).

We now turn to the RHS of the Einstein field equation (24). From equation (3), we have:

$$\bar{T}_\nu^\mu = (\bar{\rho} + \bar{P}/c^2)\bar{U}^\mu\bar{U}_\nu + \bar{P}\delta_\nu^\mu$$

where the bars, as usual, denote averaged, homogeneous background Universe quantities. Therefore, perturbing the stress-energy tensor w.r.t. the homogeneous background yields:

$$\begin{aligned}
T_\nu^\mu &= \bar{T}_\nu^\mu + \delta T_\nu^\mu \\
&= (\bar{\rho} + \delta\rho + (\bar{P} + \delta P)/c^2)\bar{U}^\mu\bar{U}_\nu + (\bar{\rho} + \bar{P}/c^2)(\delta U^\mu\bar{U}_\nu + \bar{U}^\mu\delta U_\nu) + (\bar{P} + \delta P)\delta_\nu^\mu \tag{29}
\end{aligned}$$

where we have neglected any anisotropic stress perturbation, in keeping with our approximation for the metric tensor perturbation. We can then write:

$$g_{\mu\nu}U^\mu U^\nu = (\bar{g}_{\mu\nu} + \delta g_{\mu\nu})(\bar{U}^\mu + \delta U^\mu)(\bar{U}^\nu + \delta U^\nu) = -c^2$$

and since $\bar{g}_{\mu\nu}\bar{U}^\mu\bar{U}^\nu = -c^2$ as well, this previous equation simplifies to (keeping only linear terms):

$$\delta g_{\mu\nu}\bar{U}^\mu\bar{U}^\nu + 2\bar{g}_{\mu\nu}\delta U^\mu\bar{U}^\nu = 0.$$

As $\bar{U}_\mu = -ac\delta_\mu^0$ and $\bar{U}^\mu = c/a\delta_0^\mu$, this equation then yields:

$$-2\Phi - 2a^2\delta U^0 \times \frac{c}{a} = 0, \text{ i.e. } \delta U^0 = -\frac{\Phi}{ac}.$$

Defining $\delta U^i \equiv v^i/a$, with $v^i \equiv dx^i/d\eta$ the *coordinate velocity*, we can write:

$$U^\mu = \frac{c}{a}(1 - \Phi/c^2, v^i/c),$$

and since $U_\mu = g_{\mu\nu}U^\nu$, we obtain $U_0 = g_{00}U^0 + g_{0i}U^i = -a^2(1 + 2\Phi/c^2) \times c/a(1 - \Phi/c^2)$, i.e. $U_0 = -ac(1 + \Phi/c^2)$ and $U_i = g_{i0}U^0 + g_{ij}U^j = a^2(1 - 2\Phi/c^2)\delta_{ij} \times v^j/a = av_i$, so that

$$U_\mu = -ac(1 + \Phi/c^2, -v_i/c).$$

Injecting these expressions for the 4-velocities into equation (29), we find

$$\begin{aligned} T_0^0 &= -(\bar{\rho} + \delta\rho)c^2 - \overline{(\bar{P} + \delta P)} + \overline{(\bar{\rho} + \bar{P}/c^2)(\Phi - \Phi)} + \overline{(\bar{P} + \delta P)\delta_0^0} \\ &= -(\bar{\rho} + \delta\rho)c^2 \\ T_0^i &= \overline{(\bar{\rho} + \delta\rho + (\bar{P} + \bar{P})/c^2)} \times 0 + (\bar{\rho} + \bar{P}/c^2)(-v^i c + 0) + \overline{(\bar{P} + \delta P)\delta_0^i} \\ &= -(\bar{\rho} + \bar{P}/c^2)cv^i \\ T_j^0 &= \overline{(\bar{\rho} + \delta\rho + (\bar{P} + \delta P)/c^2)} \times 0 + (\bar{\rho} + \bar{P}/c^2)(0 + v_j c) + \overline{(\bar{P} + \delta P)\delta_j^0} \\ &= (\bar{\rho} + \bar{P}/c^2)cv_j \\ T_j^i &= \overline{(\bar{\rho} + \delta\rho + (\bar{P} + \delta P)/c^2)} \times 0 + \overline{(\bar{\rho} + \bar{P}/c^2)(0 + 0)} + \overline{(\bar{P} + \delta P)\delta_j^i} \\ &= \overline{(\bar{P} + \delta P)\delta_j^i}. \end{aligned}$$

(30)

An important thing to note is that if we define the *momentum density* as $q^i \equiv (\bar{\rho} + \bar{P}/c^2)v^i$, and consider the stress energy tensor as the sum of its various component, that is to say we write

$$T_{\mu\nu} = \sum_X T_{\mu\nu X},$$

where X stands for photons, baryons, dark matter, etc ..., then the perturbations in density, pressure and momentum density of these components simply add, i.e.:

$$\delta\rho = \sum_X \delta\rho_X ; \quad \delta P = \sum_X \delta P_X ; \quad q^i = \sum_X q_X^i .$$

We can now plug all these expressions for the perturbed stress energy tensor components into equations ((26), (27) and (28)) to obtain the linearised Einstein field equations:

$$\begin{aligned} G_{00} &= -\frac{8\pi G}{c^4} g_{0\mu} T_0^\mu \\ -3\frac{\mathcal{H}^2}{c^2} - 2\frac{\nabla^2\Phi}{c^2} + 6\mathcal{H}\frac{\Phi'}{c^4} &= -\frac{8\pi G a^2}{c^2} \bar{\rho} \left(1 + \delta + 2\frac{\Phi}{c^2} \right) \end{aligned}$$

where $\delta \equiv \delta\rho/\bar{\rho}$ as usual, the 0th order terms on both sides yield the FL equation (5) in conformal coordinates for a flat spacetime and without a cosmological constant term:

$$\mathcal{H}^2 = \frac{8\pi G a^2}{3} \bar{\rho}, \tag{31}$$

and the first order ones give the 00 component of the perturbed equations (24) we are looking for:

$$\nabla^2\Phi = 4\pi G a^2 \bar{\rho} \delta + \frac{8\pi G a^2}{c^2} \bar{\rho} \Phi + 3\mathcal{H} \frac{\Phi'}{c^2} .$$

Upon substitution of equation (31) in the second RHS term, this first order equation simplifies to:

$$\nabla^2\Phi = 4\pi G a^2 \bar{\rho} \delta + 3\frac{\mathcal{H}}{c^2} (\mathcal{H}\Phi + \Phi') \tag{32}$$

which, for $c \rightarrow \infty$ gives us the Newton-Poisson equation (19). We can also see that if we assume $\Phi' \sim \mathcal{H}\Phi$ and recall that $\mathcal{H} \sim 1/\eta$, the relativistic correction term on the RHS is small in front of the LHS term when $kc\eta \gg 1$ ¹, that is when the wavelengths of the perturbations are much smaller than the particle horizon.

The ‘‘relativistic’’ Newton-Poisson equation (32) can be written in another form if we consider the mixed term linearised Einstein equation:

$$\begin{aligned} G_{0i} &= -\frac{8\pi G}{c^4} g_{0\mu} T_i^\mu \\ -\frac{2}{c^3} \partial_i (\mathcal{H}\Phi + \Phi') &= \frac{8\pi G a^2}{c^3} (\bar{\rho} + \bar{P}/c^2) v_i \end{aligned}$$

¹Obtained by taking the Fourier transform of the equation.

and integrate it assuming perturbations vanish at spatial infinity and that $v_i = \partial_i \Theta$ i.e. only the scalar part of the perturbation matters**, which gives

$$\mathcal{H}\Phi + \Phi' = -4\pi G a^2 (\bar{\rho} + \bar{P}/c^2) \Theta.$$

Injecting this expression in the second term on the RHS of equation (32) then leads to the desired form:

$$\nabla^2 \Phi = 4\pi G a^2 \bar{\rho} \left(\delta - 3 \frac{\mathcal{H}}{c^2} \left(1 + \frac{\bar{P}}{\bar{\rho} c^2} \right) \Theta \right) \equiv 4\pi G a^2 \bar{\rho} \Delta \quad (33)$$

which also is a Poisson equation but one where the gravitational field is sourced by the gauge invariant scalar quantity Δ rather than the energy density contrast δ only. Note that, once again, when $c \rightarrow \infty$, we do recover the Newtonian limit given by equation (19) as $\Delta \rightarrow \delta$.

Finally the last equation comes from the spatial component of the linearised Einstein equation:

$$G_{ij} = -\frac{8\pi G}{c^4} g_{i\mu} T_j^\mu$$

$$2 \frac{\mathcal{H}'}{c^2} + \frac{\mathcal{H}^2}{c^2} - 8\mathcal{H}' \frac{\Phi}{c^4} - 6\mathcal{H} \frac{\Phi'}{c^4} - 2 \frac{\Phi''}{c^4} - 4\mathcal{H}^2 \frac{\Phi}{c^4} = -\frac{8\pi G a^2}{c^4} \left(\bar{P} + \delta P - 2 \frac{\Phi}{c^2} \bar{P} \right).$$

In a similar way to the time component, this equation yields a 0th order part:

$$\mathcal{H}' = -\frac{4\pi G a^2}{3} \left(\bar{\rho} + 3 \frac{\bar{P}}{c^2} \right), \quad (34)$$

where we have used equation (31) to replace the \mathcal{H}^2 term on the LHS. This simply is the second FL equation (6) in conformal coordinates and without the cosmological constant term on the RHS. The linear part is thus the only new equation we have derived:

$$8\mathcal{H}' \frac{\Phi}{c^4} + 6\mathcal{H} \frac{\Phi'}{c^4} + 2 \frac{\Phi''}{c^4} + 4\mathcal{H}^2 \frac{\Phi}{c^4} = \frac{8\pi G a^2}{c^4} \left(\delta P - 2 \frac{\Phi}{c^2} \bar{P} \right).$$

Once again, we can use the 0th order equation to eliminate the \bar{P} term on the RHS, and introduce the adiabatic sound speed of the fluid $c_s^2 = \delta P / \delta \rho$, to simplify this equation a bit:

$$\Phi'' + 3\mathcal{H}\Phi' + (2\mathcal{H}' + \mathcal{H}^2)\Phi = 4\pi G a^2 c_s^2 \bar{\rho} \delta \quad (35)$$

**More specifically, this means we apply the scalar-vector-tensor decomposition to the perturbed velocity (analogous to a Helmholtz decomposition of the vector into a curl free and a divergence free part, and writing the curl free component as the gradient of a scalar – sometimes called the velocity potential), thus writing $v_i = \partial_i \Theta + \hat{v}_i$, and neglecting the vector part, \hat{v}_i , in keeping with the assumptions we made for the metric perturbations.

Although in principle the Einstein field equations (along with the equation of state of the fluid) provide a complete description of the dynamics of the system, to better make the link with the Newtonian perturbation theory developed in the previous section, we now derive the relativistic equivalent of energy conservation and the Euler equations. These two sets of equations are related through the Bianchi identities as you have seen in the third year course. We thus write the conservation of the energy-momentum tensor, $\nabla_\mu T^\mu_\nu = 0$, i.e.:

$$\partial_\mu T^\mu_\nu + \Gamma^\mu_{\mu\alpha} T^\alpha_\nu - \Gamma^\alpha_{\mu\nu} T^\mu_\alpha = 0. \quad (36)$$

Let's consider the $\nu = 0$ component of this equation first. It writes:

$$\partial_0 T^0_0 + \partial_i T^i_0 + \Gamma^0_{00} T^0_0 + \Gamma^i_{i0} T^0_0 + \Gamma^0_{0i} T^i_0 + \Gamma^j_{ji} T^i_0 - \Gamma^0_{00} T^0_0 - \Gamma^i_{00} T^0_i - \Gamma^0_{i0} T^i_0 - \Gamma^j_{i0} T^i_j = 0,$$

and injecting the expressions previously obtained for the linear connection coefficients and for the perturbed stress energy tensor, we get:

$$-\partial_0(\bar{\rho} + \delta\rho)c^2 - (\bar{\rho} + \bar{P}/c^2)c\partial_i v^i - (\mathcal{H}/c - \Phi'/c^3)\delta^i_i(\bar{\rho} + \delta\rho)c^2 + 3\partial_i\Phi(\bar{\rho} + \bar{P}/c^2)v^i/c - \partial^i\Phi(\bar{\rho} + \bar{P}/c^2)v_i/c - (\mathcal{H}/c - \Phi'/c^3)\delta^j_i(\bar{P} + \delta P)\delta^i_j = 0,$$

and keeping only up to the linear terms, this simplifies to:

$$\bar{\rho}' + \bar{\rho}'\delta + \bar{\rho}\delta' + (\bar{\rho} + \bar{P}/c^2)\frac{\partial\vec{v}}{\partial\vec{x}} + 3(\mathcal{H} - \Phi'/c^2)(\bar{\rho} + \bar{P}/c^2) + 3\mathcal{H}(\delta\rho + \delta P/c^2) = 0.$$

Splitting this equation in a 0th order and a first order one, as usual, yields:

$$\bar{\rho}' + 3\mathcal{H}\bar{\rho}\left(1 + \frac{\bar{P}}{\bar{\rho}c^2}\right) = 0,$$

which is simply the conservation of energy in the homogenous background^{††}, and

$$\delta' + \left(1 + \frac{\bar{P}}{\bar{\rho}c^2}\right)\left(\frac{\partial\vec{v}}{\partial\vec{x}} - 3\frac{\Phi'}{c^2}\right) + 3\mathcal{H}\left(\frac{c_s^2}{c^2} - \frac{\bar{P}}{\bar{\rho}c^2}\right)\delta = 0 \quad (37)$$

where we have used the 0th order equation to replace $\bar{\rho}'$. Note that this gives equation (17)^{‡‡}, in the Newtonian limit where $c \rightarrow \infty$. Once again general relativistic correction terms are small for $kc\eta \gg 1$, i.e. sub-horizon scale perturbations.

We now turn to the $\nu = i$ component of the stress-energy tensor conservation equation (36). This reads:

$$\partial_0 T^0_i + \partial_j T^j_i + \Gamma^0_{00} T^0_i + \Gamma^j_{j0} T^0_i + \Gamma^0_{0j} T^j_i + \Gamma^j_{jk} T^k_i - \Gamma^0_{0i} T^0_0 - \Gamma^j_{0i} T^0_j - \Gamma^0_{ji} T^j_0 - \Gamma^j_{ki} T^k_j = 0,$$

^{††}Obtained by differentiating equation (5) w.r.t. time (setting $\Lambda = 0$ first), and using equation (6) to get rid of the second order derivative of the expansion factor w.r.t. time.

^{‡‡}In conformal coordinates.

which yields, upon injecting the linear expressions previously calculated for all these terms,

$$\begin{aligned}
& \partial_0 ((\bar{\rho} + \bar{P}/c^2) cv_i) + \partial_j (\bar{P} + \delta P) \delta_i^j + (\mathcal{H}/c + \Phi'/c^3) (\bar{\rho} + \bar{P}/c^2) cv_i \\
& + 3(\mathcal{H}/c - \Phi'/c^3) (\bar{\rho} + \bar{P}/c^2) cv_i + \partial_j \Phi (\bar{P} + \delta P) \delta_i^j / c^2 - 3 \partial_k \Phi (\bar{P} + \delta P) \delta_i^k / c^2 \\
& + \partial_i \Phi (\bar{\rho} + \delta \rho) - (\mathcal{H}/c - \Phi'/c^3) \delta_i^j (\bar{\rho} + \bar{P}/c^2) cv_j \\
& + (\mathcal{H}/c - 4\mathcal{H}\Phi/c^3 - \Phi'/c^3) \delta_{ji} (\bar{\rho} + \bar{P}/c^2) cv^j \\
& + (\delta_i^j \partial_k \Phi + \delta_k^j \partial_i \Phi - \delta_{ki} \delta^{jl} \partial_l \Phi) (\bar{P} + \delta P) \delta_j^k / c^2 = 0.
\end{aligned}$$

Keeping only the linear terms in small quantities, we can simplify the previous expression as:

$$(\bar{\rho} + \bar{P}/c^2) v_i' + \partial_i (\delta P) + (\bar{\rho}' + \bar{P}'/c^2 + 4\mathcal{H}(\bar{\rho} + \bar{P}/c^2)) v_i + (\bar{\rho} + \bar{P}/c^2) \partial_i \Phi = 0,$$

and using once more the 0th order conservation of energy equation to re-arrange the $\bar{\rho}'$ terms, we arrive at the following, more familiar form:

$$\boxed{\vec{v}' + \mathcal{H}\vec{v} - 3\mathcal{H}\frac{c_s^2}{c^2}\vec{v} = -\frac{\partial\Phi}{\partial\vec{x}} - \frac{c_s^2}{1 + \bar{P}/(\bar{\rho}c^2)}\frac{\partial\delta}{\partial\vec{x}}} \quad (38)$$

where we have taken advantage that for adiabatic fluctuations $\bar{P}'/\bar{\rho}' = \delta P/\delta\rho = c_s^2$. When $c \rightarrow \infty$ this gives the Newtonian equation (18), but more specifically, the GR correction terms are small when the sound speed of the fluid is small compared to the speed of light.

As in the Newtonian case, taking the spatial gradient of the perturbed ‘‘relativistic’’ Euler equation (38) and using equations (37) and (33) to get rid of $\nabla\vec{v}$ and $\nabla^2\Phi$ respectively, yields a (somewhat complicated and hence not generally useful) 2nd order linear partial differential equation for δ . It does, however, simplify to yield equation (22) in the Newtonian limit, i.e. when the fluid considered is non-relativistic and the scale of the perturbation is small compared to the particle horizon, approximations generally accurate enough to describe the evolution of large scale structure. For instance, the largest super clusters of galaxies that we observe today on the sky, like the local Laniakea or the more distant Saraswati super cluster, have sizes ~ 150 Mpc, which are much smaller than the current particle horizon ($R_h(z=0) \sim 14.4$ Gpc).

5 The Evolution of Large Scale Structure

Let us therefore go back and have a look at equation (22). We can identify a number of features in the evolution of δ_k without actually solving this equation. For a start, it is quite clearly the equation of a damped harmonic oscillator with time dependent damping coefficient and spring constant. The damping (second term on the LHS) is due to the expansion of the Universe and will tend to suppress growth. The spring constant (third term in between brackets on the LHS) will change sign depending on whether k is large or small. If the positive part of the spring constant, $c_s^2 k^2$, dominates then we

should expect oscillatory behaviour in the form of acoustic waves in the fluid. If the negative term, $4\pi G a^2 \bar{\rho}$ dominates, then the evolution will be unstable and we should expect $\delta_{\mathbf{k}}$ to grow. The *physical* wavelength λ_J^* , that defines the transition between these two behaviours is given by:

$$\lambda_J = \frac{2\pi}{k_J} = c_s \left(\frac{\pi}{G\bar{\rho}} \right)^{\frac{1}{2}}$$

and is known as the *Jeans wavelength*. For $\lambda > \lambda_J$ gravitational collapse dominates and perturbations grow. For $\lambda < \lambda_J$ pressure wins and perturbations do not grow. We can have a rough idea of how a given system of particles will behave if we note that $c_s^2 \sim k_B T/m$ where k_B is the Boltzmann constant, T is the temperature and m is the mass of the individual particles. We can then rewrite the Jean length as $\lambda_J = \sqrt{\pi k_B T / (Gm\bar{\rho})}$. It is clear that a system which is hot and/or made up of light particles will have a large λ_J ; a cold system with heavy particles will have a small λ_J . It is now useful to find solutions for specific scenarios.

5.1 Pressureless Fluid in the Matter Dominated Era

In this situation we have $c_s^2 \simeq 0$ and hence $\lambda \gg \lambda_J$. We can therefore discard the term which depends on pressure in equation (22). As we further restrict ourselves to the matter dominated era, we then neglect all contributions to the energy densities of the Universe in front of that of matter, so that $\bar{\rho} \simeq \rho_M$. Working in conformal coordinates, equation (7) thus reduces to $\Omega_M[\eta] = 1$, i.e. $\rho_M[\eta] = \rho_c[\eta] = 3\mathcal{H}^2[\eta]/(8\pi a^2 G)$. In that case, the fluctuations in the pressureless fluid are also those of the component driving the expansion of the background, and equation (22) simplifies to:

$$\delta_{\mathbf{k}}'' + \mathcal{H}\delta_{\mathbf{k}}' - \frac{3}{2}\mathcal{H}^2\delta_{\mathbf{k}} = 0.$$

Finally, equation (8) yields[†] $a \propto \eta^2$, so that $\mathcal{H} = 2/\eta$ and trying power law solutions of the form $\delta_{\mathbf{k}} \propto \eta^\alpha$, one easily finds:

$$\delta_{\mathbf{k}} = C_1\eta^2 + C_2\eta^{-3} \tag{39}$$

where C_1 and C_2 are constants of integration. Rewriting both terms of this solution as a function of the expansion factor a , one gets $\delta_{\mathbf{k}} \propto a$ and $\delta_{\mathbf{k}} \propto a^{-3/2}$ respectively. As a increases with time when the Universe expands, one readily deduces that the second term decays and becomes sub-dominant very fast. We are then left with the first term which continues to grow under the effect of gravity as the Universe expands: for this reason it is generally called the *growing mode*. Note that we could have performed the same calculation using the 'normal' time t rather than the conformal time η , i.e. we could have solved equation (21) instead of (22). We would then have obtained $\delta_{\mathbf{k}} \propto t^{2/3}$ and $\delta_{\mathbf{k}} \propto t^{-1}$ for the growing and decaying mode respectively.

^{*}as opposed to the *comoving* wavelength, $\lambda_J^c = \lambda_J/a$.

[†]when recast in conformal coordinates, and after separating variables a and η and integrating.

5.2 Pressureless Matter in the Λ Dominated Era

Proceeding the same way as previously, we still have $c_s^2 \simeq 0$ and $\lambda \gg \lambda_J$ and can therefore discard the term which depends on pressure in equation (22). However, since we are in the Λ dominated era, we now neglect all contributions to the energy densities of the Universe in front of that of the cosmological constant, so that $\bar{\rho} \simeq \rho_V$. Working in conformal coordinates once again, equation (7) reduces this time to $\Omega_V[\eta] = 1$, i.e. $\rho_V[\eta] = \rho_c[\eta] = 3\mathcal{H}^2[\eta]/(8\pi a^2 G)$. As there exists by definition no fluctuation in the cosmological constant which could source perturbations in the gravitational potential[‡], equation (22) simplifies to:

$$\delta_{\mathbf{k}}'' + \mathcal{H}\delta_{\mathbf{k}}' - 4\pi G a^2 \rho_M \delta_{\mathbf{k}} = \delta_{\mathbf{k}}'' + \mathcal{H}\delta_{\mathbf{k}}' - \frac{3}{2} \frac{\mathcal{H}^2 \rho_M}{\rho_V} \delta_{\mathbf{k}} = 0.$$

where the last term has been crossed out because it is a second order effect: the pressureless fluid is *not* driving the expansion of the background Universe and as such its average density $\rho_M \ll \rho_V \simeq \bar{\rho}$. Going back to equation (8) yields $a \propto 1/\eta$, so that $\mathcal{H} = -1/\eta$ and trying once again power law solutions of the form $\delta_{\mathbf{k}} \propto \eta^\alpha$, one easily finds:

$$\delta_{\mathbf{k}} = C_1 + C_2 \eta^2 \tag{40}$$

where C_1 and C_2 are constants of integration as usual. Rewriting both terms of this solution as a function of the expansion factor a , one gets $\delta_{\mathbf{k}} = C^{\text{st}}$ and $\delta_{\mathbf{k}} \propto a^{-2}$ for each of these. Again, since a increases with time when the Universe expands, one readily deduces that the second term decays and becomes sub-dominant very fast. The first term, however, does *not* grow under the effect of gravity as was the case in the matter dominated era: it stays constant. Note that had we performed the same calculation using the 'normal' time t rather than the conformal time η , we would then have obtained $\delta_{\mathbf{k}} = C^{\text{st}}$ and $\delta_{\mathbf{k}} \propto \exp(-2\sqrt{\Lambda/3}t)$ for the (non) growing and decaying mode respectively. Clearly a faster rate of expansion[§] inhibits the growth of fluctuations.

5.3 Relativistic Fluid in the Radiation Dominated Era

The characteristic properties of the fluid will also affect how it evolves under gravity in an expanding universe. Consider the growth of perturbations of a relativistic fluid with pressure. An example of this scenario is of radiation interacting strongly with baryons before recombination. During this epoch, baryons are dissociated into protons and electrons which interact with photons through Thomson scattering. The net result is that radiation behaves as a gravitating fluid with pressure $\bar{P} = 1/3 \bar{\rho} c^2$, i.e. the adiabatic sound speed is $c_s = 1/\sqrt{3} c$. Although we could extend equations (17), (18) and

[‡]i.e. a term of the form $4\pi G \rho_V \delta_V$ with $\delta_V \neq 0$ which would then appear on the RHS of the perturbed Newton-Poisson equation (19).

[§]It is easy to show from equation (8) that the expansion factor, a , grows exponentially with time as $\exp(\sqrt{\Lambda/3}t)$ in the Λ dominated era, rather than as a power law.

(19) so that they describe a relativistic fluid[¶], and use these modified equations to derive the equivalent of equation (22), it is much more satisfactory to go to the full general relativistic equations that we have worked quite hard to derive in the previous section. We will thus follow this route and leave the exploration of the ‘modified Newtonian path’ as an exercise. In the radiation dominated era, we have $\bar{\rho} \simeq \rho_\gamma$, and switching to conformal coordinates, equation (7) reduces to $\Omega_\gamma[\eta] = 1$, i.e. $\rho_\gamma[\eta] = \rho_c[\eta] = 3\mathcal{H}^2[\eta]/(8\pi a^2 G)$. Equation (8) further yields $a \propto \eta$, so that $\mathcal{H} = 1/\eta$ and finally equations (31) and (34) give $\mathcal{H}^2 = -\mathcal{H}' = 1/\eta^2$ for the background Universe in that case.

Our starting point for the perturbations will be Einstein’s field equation (35):

$$\Phi'' + 3\mathcal{H}\Phi' + (2\mathcal{H}' + \mathcal{H}^2)\Phi - 4\pi G a^2 \bar{\rho} \delta c_s^2 = 0.$$

Re-writing the relativistic Poisson equation (32) as:

$$4\pi G a^2 \bar{\rho} \delta = \nabla^2 \Phi - 3 \frac{\mathcal{H}}{c^2} (\mathcal{H}\Phi + \Phi'), \quad (41)$$

we can use it to get rid of the δ term which appears in the previous field equation to get, upon taking the Fourier transform and using $\mathcal{H}^2 = -\mathcal{H}'$,

$$\Phi_{\mathbf{k}}'' + 4\mathcal{H}\Phi_{\mathbf{k}}' + c_s^2 k^2 \Phi_{\mathbf{k}} = 0.$$

Finally, using the expansion factor scaling with conformal time we can simplify this equation a bit further as:

$$\Phi_{\mathbf{k}}'' + \frac{4}{\eta}\Phi_{\mathbf{k}}' + \frac{c^2 k^2}{3}\Phi_{\mathbf{k}} = 0.$$

The solutions of this differential equation are:

$$\Phi_{\mathbf{k}} = A_1 \frac{\sqrt{3}}{kc\eta} j_1\left(\frac{kc\eta}{\sqrt{3}}\right) + A_2 \frac{\sqrt{3}}{kc\eta} y_1\left(\frac{kc\eta}{\sqrt{3}}\right),$$

where A_1 and A_2 are constants, $j_1(x) = \sin(x)/x^2 - \cos(x)/x$ and $y_1(x) = -\cos(x)/x^2 - \sin(x)/x$ are spherical Bessel functions of the first and second kind respectively. Considering the ‘early times’ limit, i.e. $\eta \rightarrow 0$ we deduce that the constant A_2 must vanish, as otherwise the gravitational potential would blow up since $\lim_{x \rightarrow 0} y_1(x)/x = -\infty$. Note that, contrary to what would have happened had we followed the modified Newtonian road, this solution is valid for *any* wavenumber k , even those

[¶]Basically by treating the fluid as a special relativistic fluid which sources the Newtonian gravitational potential. The perturbed equations then become:

$$\begin{aligned} \dot{\delta} + 4\nabla\vec{v}/(3a) &= 0 \\ \dot{\vec{v}} + H\vec{v} &= -\nabla\Phi/a - c^2\nabla\delta/(3a) \\ \nabla^2\Phi &= 8\pi G\bar{\rho}a^2\delta \end{aligned}$$

where both the extra third in the continuity equation and the factor 2 in the Poisson equation come from the contribution of the relativistic fluid pressure (respectively \bar{P} and $3\bar{P}$ for these two equations) to the energy density.

corresponding to super-horizon wavelengths. However, in keeping with the previous sub-sections, if we only consider sub-horizon modes, i.e. $kc\eta \gg 1$, we get the solution

$$\Phi_{\mathbf{k}} \simeq -\frac{3A_1}{(kc\eta)^2} \cos\left(\frac{kc\eta}{\sqrt{3}}\right),$$

which means that the gravitational potential on small scales oscillates at a frequency $kc/\sqrt{3}$ with an amplitude that decays with (conformal) time as η^{-2} , or in terms of expansion factor as a^{-2} .

Turning back to the energy density fluctuations in the relativistic fluid, equation (41) can be conveniently re-arranged into:

$$\delta_{\mathbf{k}} = \frac{2}{3\mathcal{H}^2} \left(-k^2\Phi_{\mathbf{k}} - 3\frac{\mathcal{H}}{c^2}(\mathcal{H}\Phi_{\mathbf{k}} + \Phi'_{\mathbf{k}}) \right) = -\left(\frac{2}{3}k^2\eta^2 + \frac{2}{c^2} \right) \Phi_{\mathbf{k}} - \frac{2\eta}{c^2} \Phi'_{\mathbf{k}}. \quad (42)$$

For sub-horizon energy density perturbations, i.e. such that $kc\eta \gg 1$, the first RHS term dominates over the other two. Plugging in the solution previously obtained for the gravitational potential, we then easily derive that:

$$\delta_{\mathbf{k}} \simeq -\frac{2}{3}k^2\eta^2\Phi_{\mathbf{k}} = \frac{2A_1}{c^2} \cos\left(\frac{kc\eta}{\sqrt{3}}\right).$$

In other words, sub-horizon energy density perturbations in the relativistic fluid oscillate around $\delta_{\mathbf{k}} = 0$ at a frequency $kc/\sqrt{3}$ with constant amplitude. Note that we could also have plugged in the full solution for the gravitational potential in terms of Bessel functions and considered all the terms in equation (42), which would allow us to conclude that for super-horizon perturbations with $kc\eta \ll 1$, both $\Phi_{\mathbf{k}}$ and $\delta_{\mathbf{k}}$ are constant. Finally let us stress that this way of deriving the evolution of perturbations which is valid on *all* scales can also be applied to the matter dominated Universe that we studied earlier on. We leave it as an exercise to show that in that case, one recovers the evolution given by equation (39) for $\delta_{\mathbf{k}}$ and that the gravitational potential $\Phi_{\mathbf{k}} = A_1 + A_2\eta^{-5}$ stays \sim constant.

5.4 Pressureless Matter in the Radiation Dominated Era

Now that we have obtained the evolution of energy density perturbations in the dominant relativistic fluid, we can look at what happens to the sub-dominant pressureless matter perturbations ($c_s^2 \simeq 0$) in the radiation dominated era, just as we did in the Λ dominated era. A notable example is cold dark matter density fluctuations during the radiation dominated era: they are decoupled from the baryon + radiation fluid. In this situation, the pressureless matter will not only play a sub-dominant role in the expansion of the Universe, i.e. $\rho_{\gamma}[\eta] \simeq \bar{\rho} = 3\mathcal{H}^2[\eta]/(8\pi a^2 G)$ will still hold, but it will also make a negligible contribution to the general relativistic version of the Newton-Poisson equation (32). In other words, on sub-horizon scales ($kc\eta \gg 1$) we will have:

$$\nabla^2\Phi = 4\pi G(\rho_{\gamma}\delta_{\gamma} + \rho_M\delta_M) = 4\pi G\rho_{\gamma} \left(\delta_{\gamma} + \frac{\rho_M}{\rho_{\gamma}}\delta_M \right) \simeq 4\pi G\rho_{\gamma}\delta_{\gamma},$$

where we have crossed out the second order term. We have already calculated the evolution of δ_γ in the previous sub-section and found that on sub-horizon scales these perturbations do not grow, but oscillate rapidly around zero, so that their time averaged amplitude is also nil. Thus, in this case, equation (22) can be simplified to:

$$\delta_{\mathbf{k}}'' + \mathcal{H}\delta_{\mathbf{k}}' = \delta_{\mathbf{k}}'' + \frac{1}{\eta}\delta_{\mathbf{k}}' = 0.$$

The solution to this equation is $\delta_{\mathbf{k}} = C_1 \ln \eta + C_2$, or if we recast things in terms of the expansion factor, $\delta_{\mathbf{k}} = C_1 \ln a + C_2$. In other words, the interesting point here is that even though perturbations in the dominant fluid (the radiation) do *not* grow, perturbations in the sub-dominant pressureless matter component do. As it turns out, this logarithmically slow growth is of key importance as it means that once recombination is finished and baryons have decoupled from photons, baryonic density fluctuation growth will be sped up as these baryons will fall in deeper dark matter potential wells. It is this very acceleration of the baryonic collapse which allows high density contrast structures such as the galaxies we observe today to form rapidly enough.

5.5 Damping of Cosmological Perturbations

There exists situations which do not fit well into the formalism we have been using so far. These occur when there is imperfect coupling between different fluid components or when the system cannot be described purely in terms of a fluid and one must revert to a more microscopic description in terms of the particle distribution function.

5.5.1 The Boltzmann Equation

How to treat the matter distribution without assuming that it is a fluid? Consider the phase space distribution function $f(\vec{x}, \vec{p})$ where \vec{x} is the position and \vec{p} is the momentum of individual particles. The evolution of f is described by the Boltzmann equation:

$$\left(\frac{\partial}{\partial t} + \frac{\partial \vec{x}}{\partial t} \cdot \nabla_{\vec{x}} + \frac{\partial \vec{p}}{\partial t} \cdot \nabla_{\vec{p}} \right) f = \mathcal{Q}(f, f)$$

where the collision term on the RHS is generally a complicated multiple integral over momentum space involving differential cross-sections. Its general relativistic generalisation writes:

$$\left(p^\alpha \frac{\partial}{\partial x^\alpha} - \Gamma_{\mu\nu}^\alpha p^\mu p^\nu \frac{\partial}{\partial p^\alpha} \right) f = \left(\frac{\partial f}{\partial t} \right)_{\text{coll}}$$

5.5.2 Damping During Recombination

Before recombination, photons and charged particles are tightly coupled together, i.e. the mean free path of the photons is negligible and this radiation and baryon combination can be treated as a single

relativistic fluid as we did in section 5.3. However as the Universe expands, the density and temperature of such a fluid drop, and photons end up not possessing enough energy to prevent protons from capturing free electrons and form neutral hydrogen. Their mean free path, given by $\lambda_{\text{mfp}} \approx 1/(\sigma_T n_e)$ where n_e is the number density of free electrons and σ_T the Thompson scattering cross-section, thus progressively approaches infinity and the Universe becomes transparent to radiation. This entails that during this recombination epoch, for the brief period of time when it transitions from negligible to infinite, the mean free path of the photons becomes finite.

In other words, as the Universe recombines (around $z \simeq 1100$), n_e plummets over a redshift change of $\Delta z \simeq 80$ and because photons and electrons are not perfectly coupled at that time, the photons will be able to random walk out of overdensities as they scatter off free electrons. In doing so they will drag matter from over dense to under dense regions and therefore damp out perturbations on scales smaller than the characteristic random walk scale. In order to derive the damping scale (known as the *Silk damping* scale) rigorously, one needs to use the Boltzmann equation previously given (see e.g. [P]), but it can be derived approximately using the following route.

Consider a photon that suffers $N = c\Delta t/\lambda_{\text{mfp}}[t]$ collisions during recombination. Between two collisions it travels a comoving distance $\lambda_{\text{mfp}}/a[t]$. Since this is a random walk, this photon acquires a mean square comoving displacement $(\Delta x)^2 = N(\lambda_{\text{mfp}}/a)^2 = c\lambda_{\text{mfp}}[t]\Delta t/a^2[t]$, so that the total mean square comoving distance it travels until the time of decoupling is:

$$x^2 \equiv \int_0^{t_\star} \frac{c dt}{\sigma_T n_e[t] a^2[t]} = \int_0^{\eta_\star} \frac{c d\eta}{\sigma_T n_e[\eta] a[\eta]} = \frac{c\eta_\star}{5\sigma_T n_e[\eta_\star] a[\eta_\star]}$$

where we have used the fact that recombination occurs during the matter dominated epoch, so that $a[\eta] = a[\eta_\star](\eta/\eta_\star)^2$ and $n_e[\eta] = n_e[\eta_\star](a[\eta]/a[\eta_\star])^{-3} = n_e[\eta_\star](\eta/\eta_\star)^{-6}$. This yields a comoving Silk damping scale,

$$\lambda_S^c \equiv x = \sqrt{\frac{c\eta_\star}{5a[\eta_\star]\sigma_T n_e[\eta_\star]}}$$

using the conformal time at decoupling, η_\star . Plugging in numbers for $c\eta_\star = 262 \text{ Mpc}$, $a[\eta_\star] \simeq 1/1101 = 9.08 \times 10^{-4}$, $\sigma_T = 6.65 \times 10^{-25} \text{ cm}^2$ and $n_e[\eta_\star] = \Omega_{b,0} \rho_{c,0}/m_p \times 1101^3 \simeq 332 \text{ cm}^{-3}$ yields $\lambda_S^c \simeq 9.2 \text{ Mpc}$. Fluctuations below this scale will be washed out.

5.5.3 Massive Neutrinos During Any Era

Massive neutrinos cannot be described as a fluid- they do not interact with each other and their evolution must be studied using the Boltzmann equation. On very large scales they will tend to cluster just like matter and radiation but on small scales, they will tend to *free-stream* i.e. move relativistically from one region of space to another. This will lead an overall damping effect, wiping out structure on small scales. The damping scale will depend on their mass and is roughly given by

$$\lambda_{FS} \simeq 40 \left(\frac{30 \text{ eV}}{M_\nu} \right) \text{ Mpc}$$




Figure 1: The power spectrum for three different models of the Universe: Λ HDM, i.e. a universe with baryons, massive neutrinos and a cosmological constant; Λ BDM, i.e. a universe with only baryons and a cosmological constant; Λ CDM, i.e. a universe with baryons, cold dark matter and a cosmological constant. Note that the first two models have much less power on small scales (large k) than the last one.

6 Building models of large scale structure

Thus far we have studied the evolution of structure in a variety of scenarios and we should, by now, have a qualitative understanding of how cosmological perturbations may evolve. We now need complete this analysis by defining the initial conditions, i.e. the seeds of structure, then characterizing how perturbations of different length scales evolve and finally identifying how we should ultimately characterize large scale structure today.

Over the decades there have been a plethora of proposals for the initial conditions of structure formation. One set of possibilities is that the Universe started off in a quasi-chaotic initial state and that the thermal initial state smoothed out the large inhomogeneities leaving a residue of fluctuations when then evolved to form structure. Clearly this is not a viable proposal unless we severely modify the nature of the Universe at those early time- as we saw in the previous section, structure on very large

scales (larger than the Jean wavelength) will tend to grow under the force of gravity. Furthermore, there is a limit to how much the Universe could homogenize set by causality so it is in fact physically impossible to implement such a simple idea.

We do, however, have a proposal that tends to smooth out the Universe and that changes the causal structure of space-time. Inflation will take a microscopic patch of the Universe which is in thermal equilibrium and is well within the Jeans wavelength at that time, and expands it to macroscopic, cosmological proportions. In doing so, Inflation solves the problem of how to homogenize but also provides a mechanism for seeding structure. We expect that, due to the quantum nature of space-time and matter, that the Universe will be riven by quantum fluctuations on microscopic scales. A period of Inflationary expansion will amplify and stretch these quantum fluctuations to macroscopic scales which will be many times larger than the cosmological horizon by the time Inflation ends. As the Universe resumes its normal expansion in the radiation era, the fluctuations will seed structure in the cosmological fluid which will then evolve in the manner described in the previous sections.

The form of the initial conditions arising from Inflation have a deeply appealing feature: they will correspond to a *random field* which has a *scale invariant* gravitational potential. In this context, a random field is a three-dimensional function which can be generated through some random process; this should not come as a surprise given that the source of the fluctuations is a quantum process. And if you think about what we are trying to do, and look at the structure of the sky, you will realize that there must be an element of randomness. Our theory won't tell us if a cluster of galaxies, or a filament of galaxies or more generally an overdensity or underdensity is going to be at an exact position in space. All we can talk about is how much more probable structures of a given size are going to be relative to others. For example, we may expect to see more structure of 1 Mpc than of 100 Mpc, but we don't know exactly where they will be. Hence we talk about our density contrast, δ , or gravitational potentials being a random fields for which we can calculate their statistical properties.

We characterize a random field in much the same way we would characterize any other random process. For example we will have that the density contrast, δ satisfies

$$\langle \delta(\vec{x}) \rangle = 0$$

and that we can characterize its variance in terms of a *correlation function*, $\xi(r)$ through

$$\xi(|\vec{x} - \vec{x}'|) \equiv \langle \delta(\vec{x})\delta(\vec{x}') \rangle$$

or alternatively in terms of its *power spectrum*

$$P(k) \equiv \langle |\delta_{\vec{k}}|^2 \rangle$$

By defining $\xi(r)$ or $P(k)$ we can characterize the statistical properties of the random field* It is often useful to consider the dimensionless version of the power spectrum, the *mass variance* which is given by

$$\Delta^2(k) = \frac{k^3 P(k)}{4\pi}$$

*This is only strictly true of the random process is Gaussian. For non-Gaussian processes one has to go further and characterize such quantities as $\langle \delta(\vec{x}_1)\delta(\vec{x}_1)\delta(\vec{x}_1) \rangle$ and higher order products. It turns out the Inflation predicts that the random fields are, to a very good approximation, Gaussian.

We now need to understand what is meant by scale invariance. Let us define the average gravitational potential in a ball of radius R to be

$$\Phi(R) = \frac{1}{V_R} \int_{V_R} d^3x \Phi(\vec{x})$$

where V_R is the volume of the ball. We can define the variance of Φ on that scale to simply be

$$\sigma_R^2(\Phi) = \langle \Phi^2(R) \rangle$$

where $\langle \dots \rangle$ is an ensemble average, i.e. an average over many possible configurations of Φ . A scale invariant spectrum corresponds to a variance which is independent of R , i.e. $\sigma_R^2(\Phi) \propto \text{constant}$. It turns out that we can relate $\sigma_R^2(\Phi)$, to $\delta(t, \vec{k})$ through the Newton-Poisson equation. Indeed we have that

$$\sigma_R^2(\Phi) \simeq \frac{k^3}{4\pi} \langle |\Phi(t, \vec{k})|^2 \rangle \propto \frac{k^3}{k^4} \langle |\delta(t, \vec{k})|^2 \rangle \quad \text{with } k = \frac{2\pi}{R}$$

If it is scale invariant we then have that the power spectrum of the density fluctuations at initial time t_i , $P_i(k) \equiv \langle |\delta(t_i, \vec{k})|^2 \rangle$ has the form

$$P_i(k) \propto k$$

In practice, choosing scale invariant initial conditions for the density field corresponds to picking the amplitude of the density field[†] to be given by $|\delta(t, \vec{k})| \propto k^{1/2}$.

Having chosen a set of initial conditions we can predict what the large scale structure of the Universe for different sets of assumptions. We will do so for three model universes: a universe where matter is solely of baryons and known as the Baryonic Dark Matter (BDM) universe ; a universe which also contains pressureless dark matter and is known as the Cold Dark Matter (CDM) universe; and a universe in which matter is dominated by massive neutrinos and is known as the Hot Dark Matter (HDM) universe.

6.1 A Baryonic Dark Matter Universe

Consider the simplest scenario we can imagine with the tools we have been give: a flat Universe consisting solely of photons, atoms (i.e. Baryons) and a cosmological constant. There is a limit on how much of the Universe can be made of baryons: the abundance of light elements restricts $\Omega_b h^2 \simeq 0.024$. With our current constraints on the Hubble constant, this means that fractional energy density in baryons must be around 5% and given that we are considering a Universe with $\Omega = 1$ we must have $\Omega_\Lambda = 0.95$. In figure 1 we plot the power spectrum of such a theory and we can clearly identify the main features. On very large scales (i.e. on scales larger than the sound horizon at equality between matter and radiation) perturbations will grow until they reach the Λ era, after

[†]This is not strictly true, otherwise we would have $\langle \delta \rangle \neq 0$ but given that we are not interested in $\delta_{\vec{k}}$ today, but in $P(k)$, this prescription will suit us

which they will be constant. On scales below the sound horizon, perturbations will initially grow, then oscillate acoustically and once the Universe recombines, they will grow again until the freeze in during the Λ era. Hence we see a series of peaks and troughs on intermediate to small scales. On very small scales, i.e. on scales which are smaller than the Silk damping scale at recombination, perturbations are severely suppressed and we can see exponential damping.

Putting all the pieces together, we find that the mass variance for this theory is roughly described by

$$\begin{aligned}\Delta^2(k) &\propto k^4 \quad \text{if } k\eta_* < 1 \\ \Delta^2(k) &\propto \frac{\cos^2\left(\frac{k\eta_*}{\sqrt{3}}\right)}{k} e^{-2(k/k_S)^2} \quad \text{if } k\eta_* > 1\end{aligned}\tag{43}$$

We can also consider a Universe in which, for some reason, we have underestimated the density of baryons. If we assume that the majority of baryons are dark (in the form of dark nuggets of matters, brown dwarfs or even black holes) we can choose $\Omega_B = 0.25$ and $\Omega_\Lambda = 0.75$. In Figure 1 we plot the power spectrum of this theory which we can call the Baryonic Dark Matter scenario (or BDM for short) and we can still the gross qualitative features we identified above.

6.2 A Cold Dark Matter Universe

An interesting scenario arises if we consider a Universe in which, once again, we have radiation, baryons and Λ but now add a component of pressureless, non-relativistic matter that does not interact with the radiation. We shall call this the Cold Dark Matter scenario (or CDM) and has a qualitatively different behaviour to the BDM case. There are effectively two regimes that will define the shape of the power spectrum. On very large scales, i.e. scales such that $k\eta \ll 1$ for all times before equality between radiation and matter, the density contrast will grow as η^2 until it reaches the Λ dominated era. For scales that cross this threshold, i.e. such that $k\eta_{eq} < 1$, the density contrast will have its evolution suppressed to logarithmic growth; this suppression in growth will last between the time the wavelength of the perturbation has the same size as the sound horizon and the transition from radiation to matter domination. During the matter era, perturbations will grow again until Λ domination sets in.

A rough estimate of the mass variance, $\Delta^2(k)$ gives us

$$\begin{aligned}\Delta^2(k) &\propto k^4 \quad \text{if } k\eta_{eq} < 1 \\ \Delta^2(k) &\propto (\ln(k))^2 \quad \text{if } k\eta_{eq} > 1\end{aligned}\tag{44}$$

The overall shape can be clearly seen in Figure 1.

6.3 A Hot Dark Matter Universe

There is yet another simple model we can consider. If we now replace the pressureless matter in the CDM model by light massive neutrinos, we will have an altogether different cosmology. The




Figure 2: A survey of galaxies over the sky illustrates there are inhomogeneities spanning a wide range of scales (2dF Galaxy Redshift Survey -based at Oxford)

motivation is clear: we know that neutrinos exist and there is even tentative evidence that they may have a mass. As we saw in the previous section, neutrinos will not evolve as a fluid and will free stream while they are relativistic, exponentially damping all perturbations on small scales. The neutrinos are weakly interacting, dark (i.e. they don't interact strongly with light) and move relativistically so can be considered a "hot" component of the Universe. For these reasons, a Universe in which neutrinos make up the bulk of the energy density today is called the Hot Dark Matter scenario (or HDM). The mass variance can be roughly approximated to

$$\Delta^2(k) \propto k^4 \exp(-k\lambda_{FS}) \quad (45)$$

and is plotted in Figure 1.

7 Comparing to observations

7.1 The spatial distribution of galaxies

A typical survey of galaxies will be like the image in Figure 2

Furthermore, we have been talking about δ , a continuous field defined over all of space. But what we actually see are galaxies, bright dots in the sky. We must relate our theory, the theory of how δ

arises and evolves with our data, a catalogue of positions of galaxies in the sky. Once again let us take as our starting point, the mean density, ρ_0 and expand it in terms of the density contrast:

$$\rho = \rho_0 \left(1 + \frac{\delta\rho}{\rho}\right) = \rho_0(1 + \delta)$$

The galaxies must trace this density somehow. For example, if the density is high (i.e. where there is an overdensity) we expect to see more galaxies, more bright dots. If the density is low, we expect to see less galaxies, a void. We can make this comparison more quantitative. Take the distribution of galaxies in the Universe and lay down balls of radius λ all over. Calculate the mass contained in each ball (i.e. add up the mass in all the galaxies contained in each ball). We have that

$$M(\lambda) = \int_{S_\lambda} d^3x \rho(\vec{x})$$

where S_λ is a sphere of radius λ . We can find the average over all balls to get

$$\bar{M}(\lambda) \equiv \langle M(\lambda) \rangle = \frac{4\pi}{3} \lambda^3 \rho_0$$

Suppose we now calculate the variance

$$[\delta M(\lambda)]^2 \equiv \langle [M(\lambda) - \bar{M}(\lambda)]^2 \rangle$$

Some balls will be heavier and others will be lighter. There should be a scatter. One can show that the mass variance is given by

$$\Delta^2(k) \equiv \left[\frac{\delta M(\lambda)}{\bar{M}(\lambda)} \right]^2 \simeq \frac{k^3}{2\pi^2} P(k) \quad \text{with } k = \frac{2\pi}{\lambda}$$

So $P(k)$ gives you the fluctuations in mass of balls with a given radius. Different types of clumping will lead to different $P(k)$ and these can be compared to the clumping that we see in the distribution of galaxies.

If we were to consider a universe with baryons and radiation, as described above, we would find a power spectrum as in Figure 1.

Clearly the problem is that perturbations in the baryons are prevented from growing because of the way they link up to radiation. One solution is to have a different form of non-relativistic matter that doesn't interact with radiation. It won't feel that baryon pressure and will have its Jean's length equal to zero. Because this matter does not interact with light, it can be called "dark matter". An example of the mass variance $\Delta(k)^2$ compared to data is shown in Figure 3.




Figure 3: The mass variance, $\Delta^2(k)$ of inhomogeneities with cold dark matter compared to the real universe

7.2 Anisotropies in the Cosmic Microwave Background

We have focused on the evolution of perturbations in the density field and how we can connect them to the distribution of galaxies. There is, of course, another very important component to the Universe, which plays a significant role at early times: radiation.

Let us briefly recall the history of radiation in the Universe. At very early times the universe is highly ionized, Hydrogen is dissociated into free electrons and protons and the mean free path of photons is effectively 0. The universe is opaque. At late times, the universe is neutral, protons and electrons are in Hydrogen atoms and photons are free to propagate. The Universe is transparent. The transition from one state to another is naturally related to the binding energy of hydrogen and the evolution of the ionization fraction

$$X \equiv \frac{n_p}{n_p + n_H} \quad (46)$$

At sufficiently early times we will find that $X = 1$, i.e. the Universe is completely ionized. As it crosses a certain threshold, electrons and protons combine to form Hydrogen. This happens when the temperature of the Universe is $T \simeq 3570K$ or $0.308eV$, i.e. when it was approximately 380,000 years old, at a redshift of $z \simeq 1100$. We would naively expect this to happen at $13.6eV$. One way to think about why this isn't so is that, at a given temperature there will always be a few photons




Figure 4: The evolution of the ionization fraction as a function of redshift

with energies larger than the average temperature. This energetic photons only become unimportant at sufficiently low temperatures.

We can now reconstruct the history of a photon left over from recombination. We know that, post recombination, a photon has been travelling along a straight path from then until now. It has travelled a distance $d_* \simeq \eta_0 - \eta_*$ where η_0 (η_*) is the conformal time today (at recombination). Before recombination, the mean free path of the photon was negligible, it was effectively standing still. So we can think of recombination as the time when these photons were released to travel through space. When we look from a fixed point in the sky, we will received photons that have travelled straight towards us since η_* , the point at which they are released will map out a two dimensional sphere, of radius d_* which is known as the surface of last scatter. We can think of this light as an image of the the surface of last scatter, a photograph of a spherical slice of the universe at η_* .

Recall that the Universe was in thermal equilibrium and hence the radiation should have a black body spectrum. And given that it is approximately homogeneous on large scales, the black body will be the same in all directions, with the peak at the same temperature in all directions. But we also know that it perturbed and hence we should see deviations. These deviations will have various contributions.

For a start, we have the Stefan-Boltzman law: $\rho_\gamma = \sigma T^4$ where σ is the Stefan Boltzman constant. But we then have

$$\frac{\delta T}{T} = \frac{1}{4} \frac{\delta \rho_\gamma}{\rho_\gamma}.$$

This seems quite easy to understand: the more compressed the radiation is, the hotter it is. So points

on the surface of last scattering which have denser radiation will look hotter. This is known as the *intrinsic term*

When the photons are released at the surface of last scattering, they will collide with one last electron or proton before they propagate towards us. That proton or electron may have a peculiar velocity, \vec{v}_B . This will impart a Doppler shift on the photon, i.e. the observed temperature will be $T' = T(1 - \vec{v}_B \cdot \vec{n})$ so that

$$\frac{\delta T}{T} = -\vec{v}_B \cdot \vec{n}$$

This is known as the *Doppler term*.

There will be gravitational effects too. If the photon is caught in a gravitational well at the surface of last scatter, it will be held back, i.e. gravitationally redshifted. The bigger the well, the colder the photon becomes:

$$\frac{\delta T}{T} = -\Phi$$

This is known as the *Sachs-Wolfe term*. Finally, as the photon propagates towards us through empty space, space time is changing and warping as it evolves. The photon will be redshifted or blue shifted according to:

$$\frac{\delta T}{T} = -2 \int_{\eta_*}^{\eta_0} d\eta \dot{\Phi} \quad (47)$$

This is known as the *Integrated Sachs-Wolfe term*.

Let us put all this together now. Recall: the surface of last scatter is a sphere with radius d_* . Suppose we look in a given direction \hat{n} . We will see

$$\frac{\delta T}{T}(\hat{n}) = \frac{1}{4} \delta(\eta_*, d_* \hat{n}) - [\vec{v}_B \cdot \hat{n}](\eta_*, d_* \hat{n}) - \Phi(\eta_*, d_* \hat{n}) - 2 \int_{\eta_*}^{\eta_0} d\eta' \dot{\Phi}[\eta', (d_* - \eta') \hat{n}] \quad (48)$$

In summary, a measurement of $\frac{\delta T}{T}$ is a snapshot of the universe at t_* , it is related to quantities that we know from studying large scale structure and we can use it to do “archeology” of the Universe.

What do we expect to see? First of all, recall that we looked at how perturbations in the radiation and baryons evolve in the radiation era, before recombination. On small scales, there should be a series of acoustic oscillations. We found a solution, $\delta_\gamma \propto J_2(k\eta/\sqrt{3})$. We also know that $\vec{\nabla} \cdot \vec{v} = -\dot{\delta}$. So if $\delta_\gamma \propto \cos(k\eta/\sqrt{3})$ then $\vec{v}_B \cdot \hat{n} \propto \sin(k\eta/\sqrt{3})$. We should see an oscillatory pattern in $\frac{\delta T}{T}$ with the spacing between peaks and troughs set by the angular projection of $k\eta_*/\sqrt{3}$. On very large scales we expect it to be relatively featureless.

On small scales, an altogether different phenomenon kicks in. During recombination, as the photons decouple from the baryons, they will slowly start to propagate. They will move around a little and in doing so, they will leak energy from high density regions to low density regions. The net effect is to reduce high density regions, fill in low density regions and effectively smooth out perturbations. Features on small scales will be smoothed out.




Figure 5: The best fit power spectrum of the CMB as compared to the WMAP and SPT data (Keisler 2011)

How do we analyse a map of the cosmic microwave background? We need to take the equivalent of the Fourier transform except now it is on the surface of the sphere. You will recall from mathematical methods and quantum mechanics that there is a useful basis to this in, the Spherical Harmonic functions. So we can take

$$\frac{\delta T}{T}(\hat{n}) = \sum_{\ell m} a_{\ell m} Y_{\ell m}(\hat{n}). \quad (49)$$

and plot the power spectrum

$$C_\ell = \frac{1}{2\ell + 1} \sum_m |a_{\ell m}|^2 \quad (50)$$

In Figure 5 you can see the structure of the power spectrum. There is clear evidence of the oscillatory structure as well as of the damping on very small scales and you can appreciate how remarkably well this power spectrum has been measured.