

## The KMOS Redshift One Spectroscopic Survey (KROSS)

**Richard Bower (ICC, Durham University) and Martin Bureau (University of Oxford), on behalf of the KROSS consortium**

**We present a brief overview of the first results from KROSS, a VLT KMOS guaranteed time program that will spatially resolve the dynamics, metallicity and star formation of 1000 mass and colour-selected galaxies at  $z\sim 1$ . These data will chart the formation of disk galaxies at the epoch of peak star-formation density in the universe.**

At redshift  $z\sim 1$ , the universe is roughly half its present age and very different to the present day. In particular, the volume averaged star formation density of the universe is an order of magnitude larger than at  $z=0$  (e.g. Lilly et al. 1996). Although recognisable, disk galaxies are just beginning to emerge from the chaos of the first few billion years of our universe's history, and their star formation rates, gas masses and (possibly) interaction rates are far higher than in today's quiescent galaxies. A second-generation VLT instrument, the K-band Multi-Object Spectrograph (KMOS) is a near-infrared integral-field spectrograph with a high multiplexing capability (Sharpley et al. 2013). Its 24 integral-field units (IFUs) allow to create maps of key astrophysical quantities faster than ever before, and they are particularly powerful to target the well-understood rest-frame optical wavelength range around  $z\sim 1$ . The KMOS Redshift One Spectroscopic Survey (KROSS) therefore aims to understand the aforementioned changes through a ground-breaking spatially-resolved study of a large and representative sample of galaxies at this key epoch.

### Observational Challenges of Galaxy Surveys

Within the past decade, there has been incredible progress in the study of galaxies in the redshift range  $1 < z < 2$ . A variety of techniques have been used to reliably disentangle distant galaxies from faint foreground objects, and the number of spectroscopically-confirmed galaxies in this redshift range now numbers well into the ten thousands. Rest-frame UV and optical colours, wide-area narrow-band surveys, and mid-infrared/sub-millimeter detections have all allowed observers to reliably determine galaxy stellar masses and quantify star formation rates. Although these data do not have the statistical power or quality of local surveys such as SDSS or 2dF (e.g. Abazajian et al. 2009; Colless et al. 2001), they do allow us to measure the evolution of stellar mass and to track black-hole growth (e.g. Alexander & Hickox 2012). It has also become possible to chart the dispersal of metals (e.g. Maiolino et al. 2008) and to begin to study the morphological and structural properties of galaxies as they settle onto a recognisable Hubble sequence (e.g. Swinbank et al. 2010). All these surveys have confirmed that the early universe looks drastically different from its current state, and they have allowed us to begin to infer the formation histories of typical galaxies. At the highest redshifts, massive galaxies are seen to be undergoing rapid star-formation and the Hubble sequence is not yet in place.

Advances in observational surveys have been mirrored by progress in numerical simulations. In particular, models have highlighted the importance of high gas accretion at  $1 < z < 2$ . At these redshifts, even massive galaxies are continuously fed by narrow and cool "cold-streams" of gas from the inter-galactic medium (Dekel et al. 2009; van der Voort 2011), promoting rapid but unstable star-formation (e.g. Hopkins et al. 2012). As the universe ages, these inflows become more tenuous and are eventually disrupted by outbursts from the black holes that grow at the centres of the emerging galaxies (e.g. Bower et al. 2006). By the present-day, these processes have led to the formation of a recognisable Hubble sequence. In the current generation of simulations, the emergence of the Hubble sequence arises because of the evolution of the fuelling rates of galaxies, and not because of a decline in the rate of galaxy merging (Behroozi et al. 2013).

Simulation results generally agree well with the global properties of the galaxy population, but they are intrinsically statistical in nature. With increasing surveys speeds and computational power, it is now possible to study the development of the internal properties of large samples of galaxies across these redshifts. Indeed, to enhance our empirical knowledge and refine theoretical models, the observational challenge must now be to quantitatively measure the

internal structures of intermediate and high-redshift galaxies across a range of mass and star-formation rate (SFR). In this way, we will chart the growth of galaxy disks, comparing the growth of rotation speed, disk stability and gas content against simulated galaxies. With sufficient angular resolution, we can test whether secular processes in early systems play a significant role in building the majority of the  $z = 0$  stellar mass, and we can identify the key physical mechanisms producing the global trends observed at  $z = 0$  (e.g. Forster-Schreiber et al. 2009; Swinbank et al. 2012). Equally important, these constraints are vital to determine if the numerical prescriptions developed to describe star-formation in local galaxies can equally be applied to the dense and rapidly evolving interstellar medium of gas-rich, intermediate-redshift galaxies.

## **KROSS**

Some progress has been made in building samples of galaxies at  $z=0.5-1$  using VLT/SINFONI and Keck/OSIRIS (e.g. Putsch et al. 2008), but with its 24 simultaneous IFUs KMOS is already revolutionising this field and allows us to rapidly build significantly larger and more representative galaxy samples at these redshifts. Comparing the new data to those of local optical IFU surveys such as SAMI and MANGA will reveal how the current Hubble sequence has emerged from the chaos of the early universe. A key goal of our KMOS guaranteed time program is to establish a large and well-defined statistical sample of galaxies at  $z\sim 1$ .

KROSS will create a database of 1000 spatially-resolved, mass-selected star-forming galaxies at the critical  $z\sim 1$  epoch. The sample size is large enough that the results can be divided into multiple bins of e.g. stellar mass (ranging from  $10^{9.5}$  to  $10^{10.7} M_{\odot}$ ), SFR, rotation speed and environment. Within each bin it will be possible to quantify trends between residuals from galaxy scaling relations, and to correlate deviations in SFR with velocity asymmetry and disk warping, thus quantifying the response of the nascent disks to interactions, gas flows and internal instabilities. Hence we will determine the relative roles of nature and nurture (and their imprecise boundary) in driving the evolution of the cosmic SFR density.

The first pilot observations to test the feasibility of the survey were taken during KMOS Science Verification time in July 2013, when we observed 39 galaxies at  $z=0.84$  from the HiZELS survey of the SSA22 field, for a total of 2 hours per configuration. All observations were taken in the YJ-band filter with a spectral resolution of  $R=3500$ . We detected 30 of the targeted galaxies and measured their spatially resolved dynamics, metallicity and star-formation rate (Sobral et al. 2013; Stott et al. 2014).

In P92, KROSS was allocated its first seven guaranteed nights to target the first 240 star-forming galaxies at  $z\sim 1$ . Target selection for the survey requires careful consideration. Selecting fainter galaxies makes it simple to fill all of the KMOS IFUs, but fainter (or more distant) galaxies are smaller and therefore less well spatially resolved. While asymptotic rotation curves can be recovered by exposing for longer, this impacts on survey speed. For the target selection, we thus imposed an optimal K-band limit of  $K_{AB}=22.5$ . Our aim is to obtain a mass-selected survey of the star-forming galaxy population, so we prioritise galaxies with blue colours ( $r-z < 1$ ). This is sufficient to include essentially all galaxies with significant star formation, but galaxies with redder colours are also included with a well-defined sampling rate.

## **KROSS Preliminary Results**

Each galaxy was observed for a total of 4.5 hours using ABA sequences (with one IFU from each spectrograph permanently dedicated to sky to improve the OH residual subtraction). Data reduction was carried out with SPARK (Davies et al. 2013). As we are primarily interested in emission line properties, we find the instrument has excellent stability, and our observations are limited by sky noise rather than systematic sky residuals. In total, we detected 93% of the targets (with most non-detections corresponding to faint or passive galaxies), and derived spatially-resolved velocity maps for 80% of the sample.

The main panel of Figure 1 shows a fraction of the data obtained in an observation. The right-hand panel shows the reduced two-dimensional spectra as they appear after correcting from instrumental effects and sky subtraction, and the individual spatial slices are clearly visible. Strong emission lines are evident, including  $H\alpha$ , [NII] and the [SII] doublet. Careful examination of the spectra immediately reveals spatial gradients in the line velocities (rotation) and ratios (abundance). Thumbnails on the left-hand side of Figure 1 show the corresponding velocity fields.

The power of KROSS lies in the speed with which it is possible to build up large samples of intermediate-redshift galaxies. In Figure 2, we show the dynamics of the  $\sim 200$  galaxies in our P92 sample with spatially-resolved emission lines, the location of each postage-stamp indicating the stellar mass and SFR of the target galaxy. The location of the so-called “main-sequence” of star forming galaxies at  $z\sim 1$  is illustrated by a dashed line (Karim et al. 2011). Even a cursory examination shows that the majority of the sample galaxies have disk-like kinematics, and that objects with strongly disturbed velocity fields are rare. However, the median ratio of the ionised gas rotation velocity to velocity dispersion  $v/\sigma$  (where  $\sigma$  is corrected for both instrumental broadening and beam smearing) is  $v/\sigma\sim 3$ , much lower than in local galaxies with a similar mass (where  $v/\sigma\sim 10$ ). Of course, local galaxies also have much lower specific SFRs, so these initial data already reinforce the idea that higher specific SFRs drive greater turbulence (or vice-versa). Further analysis will allow us to determine if the change in gas mass agrees with the high fuelling rates of  $z\sim 1$  disks predicted in the simulations.

By combining the rotation and velocity dispersion maps, we can measure the fraction of “rotationally-supported” systems in a mass selected sample of  $z\sim 1$  galaxies, and use the Tully-Fisher relation to probe the evolution of the mass-to-light ratio of galaxies with redshift (e.g. Miller et al. 2013). We find that 80% of our targets rotate regularly, approximately twice the disk fraction at  $z\sim 2$  from the SINS and AMAZE surveys (Shapiro et al. 2008; Contini et al. 2012). The ratio of stellar to dynamical mass suggests molecular gas fractions of 30—60%, consistent with the evolution inferred from CO measurements between  $z=0.4$  and 2 (e.g. Tacconi et al. 2013).

The good spectral resolution of KMOS also allows us to infer spatially-resolved chemical abundance gradients (via the [N II] /  $H\alpha$  ratio), that can be used to test ‘inside-out’ vs. ‘outside-in’ disk formation scenarios (Gilmore et al. 2002).

Finally, we can map the average properties of weaker spectral features by subtracting the local rotation velocities and then binning galaxies both spatially and as a function of stellar mass, SFR, etc. An example of such a stacked spectrum is shown in Figure 3, with the ambitious goal of searching for broad  $H\alpha$  emission arising from outflowing gas (so-called “super-winds”; e.g. Pettini et al. 2002). These features allow us to measure the mass and kinetic energy of the ejecta, presumably associated with star formation and/or active galactic nucleus-driven feedback, one of the critical uncertainties in galaxy formation modelling.

In summary, we have reported on progress with the guaranteed time KMOS Redshift One Spectroscopic Survey (KROSS). By measuring the spatially-resolved kinematics of  $\sim 1000$  galaxies at  $z\sim 1$ , this powerful database will provide deep insight into the different properties of galaxies when the universe was half its current age, and those of similar mass galaxies today. We will move from asking whether the detailed properties of galaxies at these epochs differ, to understanding the internal mechanics of how and why.

*The KROSS team would like to acknowledge STFC grants ST/I001573/1 and ST/H002456/1.*

## References

- Lilly et al. 1996 ApJ 460 1  
Sharples et al. 2013 ESO Messenger 151, 21  
Abazajian et al. 2009 ApJS 182 543  
Colless et al. 2001 MNRAS 328 1039  
Alexander & Hickox 2012 NewAR 56 93  
Maiolino et al. 2008 A&A 488 463  
Swinbank et al. 2010 MNRAS 405 234  
Dekel et al. 2009 Nature 457 451  
van der Voort et al. 2011 MNRAS 414 2458  
Bower et al. 2006 MNRAS 370 645  
Behroozi et al. 2013 ApJ 762 31  
Forster-Schreiber et al. 2009 ApJ 806 1364  
Swinbank et al. 2012 MNRAS 436 935  
Putsch et al. 2008 A&A 484 173  
Sobral et al. 2013 ApJ 779 139  
Stott et al. 2014 arXiv1407.1047  
Davies et al. 2013 A&A 558, 56  
Karim et al. 2011 ApJ 730 61  
Miller et al. 2013 ApJ 762 11  
Shapiro et al. 2008 ApJ 682 231  
Contini et al. 2012 A&A 539 91  
Tacconi et al. 2013 ApJ 768 74  
Gilmore et al. 2002 ApJ 574 39  
Pettini et al. 2002 Ap&SS 281 461






Figure 1. The main panel shows a fraction of the data obtained in a single observation. Strong emission lines are evident, as are the strong velocity and line-ratio gradients across the galaxies. The left-hand side thumbnails are the corresponding velocity fields.




Figure 2. First results from KROSS, our VLT/KMOS guaranteed time survey of 1000 mass and colour-selected galaxies at  $z \sim 1$ . The image shows the velocity fields of the  $\sim 200$  galaxies with spatially-resolved emission lines observed in P92, their position on the plot indicating their stellar mass and SFR. The dashed line shows the location of the so-called “main-sequence” of star forming galaxies at  $z \sim 1$ . Disk-like kinematics is seen in the majority of the sample.




Figure 3. Stacked spectrum of all KROSS galaxies from Science Verification and P92 that meet our selection criteria. This continuum and mean velocity of each galaxy spectrum were subtracted before constructing the stack, and each spectrum was normalised according to its reddening-corrected SFR before stacking. The insets show zooms around the OI 6300 and NaD 5995 absorption lines. The lower panel shows the residuals from the best-fit model (red).