

Figure 1: The angular correlation functions for colour selected subsamples of galaxies at $B \sim 24$ from a mosaic of 100 INT Tek pointings. The solid line confirms the result of ref[3] that the bluest galaxies (which correspond to the excess number counts) show clustering comparable to the old population of cluster galaxies (dotted line).



Figure 2: $V-I/R-I$ diagram showing the location of the expected main sequence, $O5V-M5V$ (crossed line). The points show the location of 1000 simulated quasars with $5.05 < z < 6.15$, (100 in each $\Delta z = 0.1$). A cut of $V - I > 3.5$ selects 100% of quasars with $z > 5.3$

References:

[1] Efstathiou, G., 1995, MNRAS, 272, L25.
[2] Roche, N., Shanks, T., Metcalfe, N., & Fong, R., 1996, MNRAS, 280, 397
[3] Landy, S.D., Szalay, A.S., & Koo, D.C., 1996, ApJ, 460, 94
[4] Dalton, G.B., Boyle, B.J., & Efstathiou, G., *in preparation*
[5] Geller, M.J., De Lapparent, V. & Kurtz, M.J., 1984, Ap.J., **287**, L55.
[6] Frieman, J.A., & Gaztañaga, E., 1994, Ap.J., **437**, L13.
[7] Schmidt, M., Schneider, D.P., & Gunn, J.E., 1991, *in: The Space Distribution of Quasars*, p.109.
[8] Warren, S.J., Hewett, P.C., & Osmer, P.S., 1994, Ap.J., **421**, 412.
[9] Hu, E., & McMahon, R.G., 1996, Nature, **382**, 281.
[10] Dalton, G.B., Maddox, S.J., Sutherland, W.J., & Efstathiou, G., 1997, MNRAS, **289**, 263.
[11] Postman, M., *et al.*, 1996, A.J., **111**, 615.
[12] Dalton, G.B., Efstathiou, G., Maddox, S.J., & Sutherland, W.J., 1994, MNRAS, **269**, 151.
[13] Dalton, G.B., *et al.*, 1994, MNRAS, 247, L47.
[14] Steidel, C.C., *et al.*, 1996, Ap.J., **462**, L17. See also astro-ph/9802318