


## Evolution of Massive Stars




Maeder (1987)


## EVOLUTION OF MASSIVE STARS ( $M \gtrsim 13 M_{\odot}$ )

- massive stars continue to burn nuclear fuel beyond hydrogen and helium burning and ultimately form an *iron core*
- alternation of nuclear *burning* and *contraction* phases

▷ *carbon burning* ( $T \sim 6 \times 10^8 \text{ K}$ )



▷ *oxygen burning* ( $T \sim 10^9 \text{ K}$ )




▷ *silicon burning*: photodisintegration of complex nuclei, hundreds of reactions  $\rightarrow$  *iron*


▷ form *iron core*

▷ *iron* is the most tightly *bound nucleus*  $\rightarrow$  no further energy from nuclear fusion

▷ iron core surrounded by *onion-like shell structure*



Helium Core (8 Msun)



Itoh and Nomoto (1987)

## EXPLOSION MECHANISMS


- two main, completely different mechanisms

### Core-Collapse Supernovae



- triggered after the exhaustion of nuclear fuel in the core of a massive star, if the *iron core mass* > *Chandrasekhar mass*
- energy source is *gravitational energy* from the collapsing core ( $\sim 10\%$  of neutron star rest mass  $\sim 3 \times 10^{46}$  J)
- most of the energy comes out in *neutrinos* (SN 1987A!)
  - ▷ *unsolved problem*: how is some of the neutrino energy *deposited* ( $\sim 1\%$ ,  $10^{44}$  J) in the envelope to *eject* the envelope and produce the supernova?
- leaves *compact remnant* (neutron star/black hole)

### Thermonuclear Explosions




- occurs in *accreting carbon/oxygen white dwarf* when it reaches the *Chandrasekhar mass*
  - *carbon ignited* under *degenerate* conditions; nuclear burning raises T, but not P
  - *thermonuclear runaway*
  - *incineration* and *complete destruction* of the star
- energy source is *nuclear energy* ( $10^{44}$  J)
- *no compact remnant* expected
- main producer of *iron*
- *standard candle* (Hubble constant, acceleration of Universe?)

but: *progenitor* evolution not understood


- ▷ *single-degenerate channel*: accretion from non-degenerate companion
- ▷ *double-degenerate channel*: merger of two CO white dwarfs

## Core Collapse




- *central properties* at the beginning of core collapse: for  $M_{\text{core}} = 1.5 M_{\odot}$ ,  $T_c \simeq 8 \times 10^9 \text{ K}$ ,  $\rho_c \simeq 4 \times 10^{12} \text{ kg m}^{-3}$
- instabilities in the contracting core lead to essentially *free-fall* collapse
- *photodissociation of nuclei*
  - ▷  $T_c \sim 10^{11} \text{ K}$  :  $\gamma + {}^{56}\text{Fe} \rightleftharpoons 13 \alpha + 4 n - 124 \text{ Mev}$
  - ▷ *endothermic reaction* (requires heat)  $\rightarrow$  temperature increases less rapidly than pressure  $\rightarrow$  rapid contraction
  - ▷  $T_c \sim 2 \times 10^{11} \text{ K}$  :  $\gamma + {}^4\text{He} \rightleftharpoons 2 p + 2 n - 28 \text{ Mev}$
- *note*: all of these reactions occur in both directions; *maximization of entropy* favours right-hand sides (larger number of particles)
- these reactions essentially undo all of the previous nuclear fusion reactions
- *neutronization*
  - ▷ *electron capture* reactions (reduce the number of electrons and electron degeneracy pressure)  $e^- + (Z, A) \rightarrow \nu_e + (Z - 1, A)$
  - $e^- + p \rightarrow \nu_e + n$  (also:  $n \rightarrow \bar{\nu}_e + p + e^-$ )
- most of the energy is lost by *neutrino emission* (10 % of the rest mass energy of the neutron star)
- *energy source*: gravitational energy

## 312 FILIPPENKO **Supernova Classification**



## SUPERNOVA CLASSIFICATION



*observational:*

- *Type I:* no hydrogen lines in spectrum
- *Type II:* hydrogen lines in spectrum

*theoretical:*

- *thermonuclear explosion* of degenerate core
- *core collapse* → neutron star/black hole


*relation no longer 1 to 1 → confusion*

- *Type Ia* (Si lines): thermonuclear explosion of white dwarf
- *Type Ib/Ic* (no Si; He or no He): core collapse of He star
- *Type II-P:* “classical” core collapse of a massive star with hydrogen envelope
- *Type II-L:* supernova with linear lightcurve (thermonuclear explosion of intermediate-mass star? probably not!)

*complications*

- special supernovae like *SN 1987A*
- *Type IIb:* supernovae that change type, *SN 1993J* (Type II → Type Ib)
- some supernova “*types*” (e.g., IIn) occur for both explosion types (“*phenomenon*”, not type; also see SNe Ic)
- new types: thermonuclear explosion of He star (Type Iab?)

# Supernova Lightcurves



- **Explosion energy:**  $E \sim 10^{44} \text{ W}$  ( $\sim$  binding energy of Fe core  $\sim GM_{\text{Fe}}^2/R_{\text{Fe}}$  with  $M_{\text{Fe}} \sim 1 M_{\odot}$ ,  $R_{\text{Fe}} \sim 2 \times 10^6 \text{ m}$ )
- much larger than the binding energy of the envelope (for  $R \sim 10^3 R_{\odot}$ )

→ **kinetic energy**  $E \sim M_{\text{env}} v^2 / 2$

$$v \simeq \left( \frac{2E}{M_{\text{env}}} \right)^{1/2} \sim 3000 \text{ km s}^{-1}$$

- energy *diffuses* out of the expanding ejecta (radius  $R$ )
- **diffusion time**,  $t_{\text{diff}}$ :  $t_{\text{diff}} \simeq R^2/(lc)$ , where the mean free path  $l$  is given by  $l = \frac{1}{\kappa\rho} \simeq \frac{4R^3}{\kappa M_{\text{env}}} \rightarrow t_{\text{diff}} \sim \frac{M_{\text{env}}\kappa}{4Rc}$
- **but:**  $R(t) \simeq vt$ , substitute and solve for  $t = t_{\text{diff}}$

$$t_{\text{diff}} \simeq \frac{M_{\text{env}}^{3/4} \kappa^{1/2}}{2(2E)^{1/4} c^{1/2}} \simeq 150 \text{ d}$$

(for  $E = 10^{44} \text{ J}$ ,  $M_{\text{env}} = 10 M_{\odot}$ ,  $\kappa = 0.034 \text{ m}^2/\text{kg}$ )

- **peak luminosity:**  $L_{\text{peak}} \sim E/t_{\text{diff}} \sim 8 \times 10^{36} \text{ W}$  (a bit high)

- late-time light curve is powered by *radioactive decay of Ni and Co*




- releasing  $5.9 \times 10^{41} \text{ J}$  and  $1.3 \times 10^{42} \text{ J}$  for each  $0.1 M_{\odot}$  of Ni

- **radioactive luminosity:**  $L(t) = L_0 e^{-t/\tau}$ , where  $\tau = t_{1/2}/\ln 2 \simeq 112 \text{ d}$

- $E_{\text{tot}} = \int_0^{\infty} L(t) dt = \tau L_0 = M_{\text{Ni}} c^2 \epsilon_{\text{Co}}$ ,  $\epsilon_{\text{Co}} \simeq 7 \times 10^{-5}$


$$\rightarrow L_{\text{radioact}} \simeq 1.3 \times 10^{35} \text{ W} \left( \frac{M_{\text{Ni}}}{0.1 M_{\odot}} \right) \exp\left( \frac{-t}{112 \text{ d}} \right)$$

## Supernova lightcurves (core collapse)



### LIGHTCURVES OF CORE-COLLAPSE SUPERNOVAE

- central explosion may be very similar in all cases (with  $E \sim 10^{44}$  J)
- *variation* of lightcurves/supernova subtypes mainly due to varying *envelope properties*
  - ▷ *envelope mass*: determines thermal diffusion time and length/existence of plateau
  - ▷ *envelope radius*: more compact progenitor → more expansion work required → dimmer supernova
- binary interactions mainly affect stellar envelopes
- a large fraction of all stars are in interacting binaries
  - *binary interactions are, at least in part, responsible for the large variety of supernova (sub-)types*
- *recent*: new-born pulsars (neutron stars) have large space velocities (median:  $200 - 300 \text{ km s}^{-1}$ )
  - neutron-stars receive a *large supernova kick*
    - ▷ probably due to *asymmetry in neutrino flux* (1%)
    - ▷ momentum balance:
 
$$M_{\text{NS}} v_{\text{kick}} = \epsilon \frac{E_{\nu}}{c} \text{ (neutrino momentum)}$$
  - $v_{\text{kick}} \simeq 350 \text{ km s}^{-1} \left( \frac{\epsilon}{0.01} \right)$   
 ( $\epsilon$ : anisotropy factor,  $M_{\text{NS}} = 1.4 M_{\odot}$ ,  $E_{\nu} = 3 \times 10^{46} \text{ J}$ )




Hsu, Ross, Joss, P.

## TYPE IA SUPERNOVAE


<http://www-supernova.lbl.gov/>

C. Panypacker  
M. DellaValle  
R. Ellis, R. McMahon  
B. Schaefer  
P. Ruiz-Lapuente  
H. Newberg  
Yale University  
Univ. of Padova  
Univ. of Barcelona  
IoA, Cambridge  
Fermilab



We have discovered well over 50 high redshift Type Ia supernovae so far. Of these, approximately 50 have been followed with spectroscopy and photometry over two months of the light curve. The redshifts shown in this histogram are color coded to show the increasing depth of the search with each new "batch" of supernova discoveries. The most recent supernovae, discovered the last week of 1997, are now being followed over their lightcurves with ground-based and (for those labeled "HST") with the Hubble Space Telescope.

### Low Redshift Type Ia Template Lightcurves



Type Ia supernovae observed "nearly" show a relationship between their peak absolute luminosity and the timescale of their light curve: the brighter supernovae are slower and the fainter supernovae are faster (see Phillips, Ap J Lett., 1993 and Riess, Press, & Kirshner, Ap J Lett., 1995). We have found that a simple linear relation between the absolute magnitude and a "stretch factor" multiplying the lightcurve timescale fits the data quite well until over 45 restframe days post peak. The lower plot shows the "nearly" supernovae from the upper plot, after fitting and removing the stretch factor, and "correcting" peak magnitude with this simple calibration relation.

- recently: Type Ia supernovae have been used as *standard distance candles* to measure the *curvature* of the Universe → *accelerating Universe?*
- Type Ia supernovae are no good standard candles! (peak luminosities vary by a factor up to 10)
- but they may be *standardizable candles*, i.e. there *appears* to be a unique relation between peak luminosity and the width of the lightcurve which can be used to derive good distances

### Caveats:

- the relation between lightcurve shape and peak luminosity is not well understood (depends on diffusion time and probably opacity)
- *the progenitors of Type Ia supernovae are not known*
- *many progenitors models*
  - ▷ *Chandrasekhar white dwarf accreting* from a companion star (main-sequence star, helium star, subgiant, giant)  
Problem: requires *fine-tuning* of accretion rate
  - ▷ *merging of two CO white dwarfs* with a total mass > Chandrasekhar mass (probably not, more likely to lead to formation of neutron star)
  - ▷ *sub-Chandrasekhar mass white dwarfs* (helium shell flash leading to a detonation of the white dwarf; extremely unlikely!)

## END STATES OF STARS

### IMPORTANT STELLAR TIMESCALES

- *dynamical timescale*:  $t_{\text{dyn}} \simeq \frac{1}{\sqrt{4G\rho}}$   
 $\sim 30 \text{ min } (\rho/1000 \text{ kg m}^{-3})^{-1/2}$
- *thermal timescale* (Kelvin-Helmholtz):  $t_{\text{KH}} \simeq \frac{GM^2}{2RL}$   
 $\sim 1.5 \times 10^7 \text{ yr } (M/M_{\odot})^2 (R/R_{\odot})^{-1} (L/L_{\odot})^{-1}$
- *nuclear timescale*:  $t_{\text{nuc}} \simeq \frac{M_c/M}{\text{efficiency}} \frac{\eta}{(Mc^2)/L}$   
 $\sim 10^{10} \text{ yr } (M/M_{\odot})^{-3}$

Three (main) possibilities

- the star develops a *degenerate core* and nuclear burning stops (+ envelope loss) → *degenerate dwarf (white dwarf)*
- the star develops a *degenerate core* and *ignites* nuclear fuel *explosively* (e.g. carbon) → *complete disruption in a supernova*
- the star *exhausts* all of its *nuclear fuel* and the core exceeds the *Chandrasekhar mass* → *core collapse, compact remnant (neutron star, black hole)*

Final fate as a function of initial mass ( $M_0$ ) for  $Z = 0.02$

Example	$t_{\text{dyn}}$	$t_{\text{KH}}$	$t_{\text{nuc}}$
<i>main-sequence stars</i>			
a) $M = 0.1 M_{\odot}$ , $L = 10^{-3} L_{\odot}$ , $R = 0.15 R_{\odot}$	4 min	$10^9 \text{ yr}$	$10^{12} \text{ yr}$
b) $M = 1 M_{\odot}$ , $L = 1 L_{\odot}$ , $R = 1 R_{\odot}$	30 min	$15 \times 10^6 \text{ yr}$	$10^{10} \text{ yr}$
c) $M = 30 M_{\odot}$ , $L = 2 \times 10^5 L_{\odot}$ , $R = 20 R_{\odot}$	400 min	$3 \times 10^3 \text{ yr}$	$2 \times 10^6 \text{ yr}$
<i>red giant</i> ( $M = 1 M_{\odot}$ , $L = 10^3 L_{\odot}$ , $R = 200 R_{\odot}$ )	50 d	75 yr	
<i>white dwarf</i> ( $M = 1 M_{\odot}$ , $L = 5 \times 10^{-3} L_{\odot}$ , $R = 2.6 \times 10^{-3} R_{\odot}$ )	7 s	$10^{11} \text{ yr}$	
<i>neutron star</i> ( $M = 1.4 M_{\odot}$ , $L = 0.2 L_{\odot}$ , $R = 10 \text{ km}$ , $T_{\text{eff}} = 10^6 \text{ K}$ )	0.1 ms	$10^{13} \text{ yr}$	

$M_0 \lesssim 0.08 M_{\odot}$	<i>no hydrogen burning</i> (degeneracy pressure + Coulomb forces)	<i>planets, brown dwarfs</i>
$[0.08, 0.48] M_{\odot}$	<i>hydrogen burning, no helium burning</i>	<i>degenerate He dwarf</i>
$[0.48, 8] M_{\odot}$	<i>hydrogen, helium burning</i>	<i>degenerate CO dwarf</i>
$[8, 13] M_{\odot}$	<i>complicated burning sequences, no iron core</i>	<i>neutron star</i>
$[13, 80] M_{\odot}$	formation of <i>iron core, core collapse</i>	<i>neutron star, black hole</i>
$M_0 \gtrsim 80 M_{\odot}$	<i>pair instability?</i> complete disruption?	<i>no remnant</i>
also (?) $[6, 8] M_{\odot}$	<i>degenerate carbon ignition possible (but unlikely), complete disruption</i>	<i>no remnant</i>